

Cathepsin S provokes interleukin-6 (IL-6) trans-signaling through cleavage of the IL-6 receptor in vitro

Charlotte M. Flynn¹, Yvonne Garbers², Stefan Düsterhöft³, Rielana Wichert¹, Juliane Lokau⁴, Christian H.K. Lehmann^{5,6}, Diana Dudziak^{5,6}, Bernd Schröder⁷, Christoph Becker-Pauly¹, Stefan Rose-John¹, Samadhi Aparicio-Siegmund¹, and Christoph Garbers^{4,*}

¹Institute of Biochemistry, Kiel University, Kiel, Germany;

²Institute of Psychology, Kiel University, Kiel, Germany;

³Institute of Molecular Pharmacology, RWTH Aachen University, Aachen, Germany;

⁴Department of Pathology, Otto-von-Guericke-University Magdeburg, Medical Faculty, Magdeburg, Germany;


⁵Laboratory of Dendritic Cell Biology, Department of Dermatology, University Hospital Erlangen, Friedrich-Alexander University of Erlangen-Nürnberg, Erlangen, Germany;

⁶Deutsches Zentrum Immuntherapie (DZI) and Medical Immunology Campus Erlangen, Germany


⁷Institute for Physiological Chemistry, Medizinisch-Theoretisches Zentrum MTZ, Technische Universität Dresden, Fiedlerstraße 42, 01307, Dresden, Germany

*Correspondence to: Dr. Christoph Garbers (christoph.garbers@med.ovgu.de, +49 391 67-15488), Department of Pathology, Otto-von-Guericke-University Magdeburg, Medical Faculty, Magdeburg, Germany; Fax: +49 391 67-15818


Figure 5a




sIL-6R



IL-6R



GAPDH




Cathepsin S




GAPDH

Figure 5c




IL-6R




IL-6R

Figure 5d



IL-6R



GAPDH