Development of Immunohistochemistry and *In Situ* Hybridisation for the Detection of SARS-CoV and SARS-CoV-2 in Formalin-Fixed Paraffin-Embedded Specimens

Fabian Z. X. Lean^{1*}, Mart M. Lamers², Samuel P. Smith^{3,7}, Rebecca Shipley^{3,6}, Debby Schipper², Nigel Temperton⁴, Bart L. Haagmans², Ashley C. Banyard³, Kevin R. Bewley⁵, Miles W. Carroll⁵, Sharon M. Brookes³, Ian Brown³, Alejandro Nuñez¹

¹ Pathology Department, Animal and Plant Health Agency (APHA), Woodham Lane, New Haw, Addlestone, KT15 3NB, United Kingdom

² Department of Viroscience, Erasmus Medical Centre, Rotterdam, Netherlands

³ Virology Department, Animal and Plant Health Agency (APHA), Woodham Lane, New Haw, Addlestone, KT15 3NB, United Kingdom

⁴ Viral Pseudotype Unit, Medway School of Pharmacy, Universities of Greenwich and Kent at Medway, Chatham ME4 4TB, United Kingdom

⁵ National Infection Service, Public Health England, Porton Down, Salisbury, SP4 0JG, United Kingdom

⁶ School of Life Sciences, University of Sussex, Falmer, Brighton, BN1 9QG, United Kingdom

⁷ Institute for Infection and Immunity, St George's Hospital Medical School, University of London, London, SW17 0RE, United Kingdom.

*Corresponding author: Fabian.lean@apha.gov.uk



Supplementary Figure 1. Immunohistochemical labelling of FFPE SARS-CoV-2 infected cells and uninfected cells. Immunodetection performed using SARS-CoV nucleoprotein rabbit polyclonal antibody (a, b) and double-stranded RNA (dsRNA) mouse monoclonal antibody (c, d) in Erasmus Medical Centre, Rotterdam. Aminoethyl carbazole (AEC) was used as the chromogen (red). Original magnification 100x.