

**Supplemental Table 1.** Association of American Urologic Association Symptom Index storage and voiding subscore\* cross-classification with frailty phenotype among 5979 men enrolled in MrOS.

<b>Multivariable Multinomial Odds Ratio (95% Confidence Interval)†</b>			
<b>Intermediate vs Robust</b>			
<b>AUASI voiding subscore</b>			
<b>AUASI storage subscore</b>	0-1	2-4	5-20
0-3	1.00	0.96 (0.79, 1.17)	0.89 (0.67, 1.18)
4-6	0.97 (0.79, 1.18)	1.03 (0.84, 1.26)	1.10 (0.91, 1.33)
7-15	1.38 (0.99, 1.92)	1.25 (0.96, 1.62)	1.18 (0.99, 1.41)
<b>Frail vs Robust</b>			
<b>AUASI voiding subscore</b>			
<b>AUASI storage subscore</b>	0-1	2-4	5-20
0-3	1.00	1.06 (0.72, 1.56)	0.95 (0.55, 1.64)
4-6	1.14 (0.77, 1.69)	1.05 (0.71, 1.56)	1.35 (0.95, 1.93)
7-15	1.88 (1.07, 3.28)	1.61 (1.03, 2.53)	2.20 (1.64, 2.96)

AUASI American Urologic Association Symptom Index

\* AUASI score range is 0 to 35 and equals the sum of 2 validated subscores based on symptom type: storage (urgency, frequency, nocturia) and voiding (intermittency, weak stream, straining, incomplete emptying). Higher score indicates more frequent symptoms. AUASI subscores categorized into approximately equal thirds (tertiles) of the subscore distribution.

† Multivariable multinomial regression model adjusted for study site, age, and self-reported history of cardiovascular disease (myocardial infarction or angina), diabetes, congestive heart failure, chronic obstructive pulmonary disease, or hypertension.