Supplemental material

BMJ Publishing Group Limited (BMJ) disclaims all liability and responsibility arising from any reliance placed on this supplemental material which has been supplied by the author(s)

BMJ Open

Supplementary Table 1: Current CVD risk factors of HDP and non-HDP women (Multiple answers collected) in order of frequency

	GH	PE	CH	Total HDP	non-HDP	P HDP vs non HDP
	n=15	n=143	n=16	n=174	n=92	
	n (%)*	n (%)*	n (%)*	n (%)*	n (%)*	
None of the options	6 (40)	52 (36)	2 (13)	60 (34)	57 (62)	<0.001
Obesity	5 (33)	38 (27)	7 (44)	50 (29)	10 (11)	0.001
Family History Heart disease	3 (20)	37 (26)	6 (38)	46 (26)	14 (15)	0.037
Hypertension	3 (20)	30 (21)	12 (75)	35 (20)	0 (0)	<0.001
Renal problems	0 (0)	21 (15)	2 (13)	23 (13)	0 (0)	<0.001
Smoking	0 (0)	10 (7)	3 (19)	13 (7)	2 (2)	0.075
History of cardiovascular event**	0 (0)	1 (1)	0 (0)	1 (1)	0 (0)	0.356
Diabetes	0 (0)	5 (3)	2 (13)	7 (4)	0 (0)	0.014
Alcohol consumption	0 (0)	0 (0)	1 (6)	1 (1)	0 (0)	0.356
High Cholesterol	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	-

*Table represents frequency of each option; percentages add to over 100% as women were asked to select any/all that applied.
**angina, stroke or heart attack
PE = preeclampsia GH = gestational hypertension CH= chronic hypertension worsening in pregnancy and/or with superimposed preeclampsia,
HDP= hypertensive disorder of pregnancy