

JVDI: Supplementary material

Stout AE, et al. Infectious disease surveillance of apparently healthy horses at a multi-day show using a novel nanoscale real-time PCR panel

**Supplementary Table 1.** Primers and probes unique to the enteric nanoscale PCR panel used in our study.

Pathogen target	Forward primer	Reverse primer	Probe	Reference
Equine rotavirus (VP4)	RGGTGGTGAYTATAMSTT CGCATT	TYACAGCGCTCCAGTCATTAC	CAGTRGGTCAGTGGC	This study
Bovine rotavirus (VP6)	TCAACATGG ATGTCCTKT AYTCCT	YCCYCCAGT TTGGAAYTCATT	TCAAAAACTCTAAAGATGCW AG	1
<i>E. coli</i> CNF1	GCAAGAGAGTGTCA GTTC TGCATT	TGCCTGATGACCTTTCTCATT	TTCAAGTCACCACCTCCAAAG GCAAA	5
<i>E. coli</i> STX2	TTTGTYACTGTSACAGCW GAAGCYTTA CG	CCCCAGTTCARWGTRAGRTCM ACRTC	TCGTCAGGC ACTGTCTGAAACT GCTCC	6
<i>E. coli</i> K99	GCTATTAGTGGTCATGGC ACTGTAG	TTTGTTCGCTAGGCAGTCAT TA	ATTTTAAACTAAAACCAGCGCC CGGCA	8
<i>E. coli</i> F17	GCAGAAAATTCAATT TAT CCTT	AATTGTACCGTCATAAGCAAGC	GGCGGCTGCGTCATCTTGCT	8
<i>E. coli</i> F41	CTGCTGATTGGACGGAAG GT	CCAGTCTCCATAGCCATTAA CAG	TGATGTAATT CACCACCAATA ATAATGT CACCTG	8
<i>E. coli</i> 987p	CCAAAGTATTCCACTGCA AGCA	GCCGTAACTCCACC GTTTGT	ACATCGGAACCACCAACAGGGA ATCCT	8
<i>E. coli</i> Sta	GCAAATCCGTTAACTA ATCTAAA	ACAGAAATAAAATTGCCAAC ATTAGC	TTACCTCCCGTCATGTTGTTCA CGGAT	8
<i>E. coli</i> LT	CCGGCAGAGGATGGTA CAG	GAATCCAGGGTTCTCTCTCCA A	TAGCAGGTTCCCACCGGATCA CC	8
<i>Clostridioides difficile</i>	GGTATTACCTAATGCTCC AAATAG	TTTGTGCCATCTTCTAAGC	CCTGGTGTCCATCCTGTTTC	4
<i>Clostridium perfringens</i>	AAATGTAACAGCAGGGG CA	TGAAATTGCAGCAACTCTAGC	ATGTTCTTTCCATTACAGG CTTAGAA	3

<i>Cryptosporidium</i> spp.	GGGTTGTATTTATTAGAT AAAGAACCA	AGGCCAATACCTACCGTCT	GTGACATATCATTCAAGTTCT GAC	7
<i>Cryptosporidium parvum</i>	CAAATTGATACCGTTGT CCTCTG	GGCATGTCGATTCTAATTCA GC	TGCCATACATTGTTGTCCTGAC AAATTGAAT	2
<i>Listeria</i> spp.	Commercial assay (MicroSEQ <i>Listeria</i> spp. detection kit)			Applied Biosystems 4427410
<i>Listeria monocytogenes</i>	Commercial assay (MicroSEQ <i>Listeria monocytogenes</i> detection Kit)			Applied Biosystems 4403874

### References

- Cho Y-I, et al. Development of a panel of multiplex real-time polymerase chain reaction assays for simultaneous detection of major agents causing calf diarrhea in feces. *J Vet Diagn Invest* 2010;22:509–517.
- Guy RA, et al. Real-time PCR for quantification of *Giardia* and *Cryptosporidium* in environmental water samples and sewage. *Appl Environ Microbiol* 2003;69:5178–5185.
- Karpowicz E, et al. qPCR quantification and genetic characterization of *Clostridium perfringens* populations in biosolids composted for 2 years. *J Appl Microbiol* 2010;108:571–581.
- Liu J, et al. A laboratory-developed TaqMan Array Card for simultaneous detection of 19 enteropathogens. *J Clin Microbiol* 2013;51:472–480.
- Massoud BZ, et al. Characterization of uropathogenic *Escherichia coli* strains isolated from community acquired and hospital acquired infections in Alexandria. *Egypt J Med Microbiol* 2007;16:513–520.

6. Perelle S, et al. Detection by 5'-nuclease PCR of Shiga-toxin producing *Escherichia coli* O26, O55, O91, O103, O111, O113, O145 and O157:H7, associated with the world's most frequent clinical cases. *Mol Cell Probes* 2004;18:185–192.
- 7 . Stroup SE, et al. Real-time PCR detection and speciation of *Cryptosporidium* infection using Scorpion probes. *J Med Microbiol* 2006;55:1217–1222.
8. West DM, et al. Rapid detection of *Escherichia coli* virulence factor genes using multiplex real-time TaqMan PCR assays. *Vet Microbiol* 2007;122:323–331.