

## Supplementary Online Content

Gaudino M, Samadashvili Z, Hameed I, Chikwe J, Girardi LN, Hannan EL. Differences in long-term outcomes after coronary artery bypass surgery using single vs multiple arterial grafts and the association with sex. *JAMA Cardiol*. Published online December 23, 2020. doi:10.1001/jamacardio.2020.6585

**eFigure.** Details of Patient Inclusion and Exclusion

**eTable 1.** Provider Volume and Baseline Patient Characteristics of Male Patients Undergoing Single-Arterial and Multiple-Arterial Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

**eTable 2.** Provider Volume and Baseline Patient Characteristics of Female Patients Undergoing Single-Arterial (SAG) and Multiple-Arterial (MAG) Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

**eTable 3.** Provider Volume and Baseline Characteristics of Propensity-Matched Male Patients Undergoing Single-Arterial and Multiple-Arterial Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

**eTable 4.** Provider Volume and Baseline Characteristics of Propensity-Matched Female Patients Undergoing Single-Arterial (SAG) and Multiple-Arterial (MAG) Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

**eTable 5.** Operative (In-Hospital/30 day) Mortality Rates of Female and Male Propensity Matched Patients

**eTable 6.** Seven-Year Mortality Rates of Propensity-Matched Risk-Stratified Female and Male Patients According to Different Low Risk Score Cut-Offs After Multiple- (MAG) vs Single-Arterial (SAG) Coronary Artery Bypass Grafting in New York State Between 2005-2014

**eTable 7.** Fully Adjusted Cox Regression Models Evaluating the Effect of Multiple Arterial Grafting (MAG) on Death in the Overall Population

**eTable 8.** Landmark Analysis of Alive Female and Male Low- and High-Risk Patients at 30 Days


**eTable 9.** Sensitivity Analysis Using Gamma Testing

**eTable 10.** Sensitivity Analysis Using E-Value

**eTable 11.** Fully Adjusted Cox Regression Models Comparing Mortality Between Female and Male Patients Receiving Single Arterial Grafts (SAG), and Female and Male Patients Receiving Multiple Arterial Grafts (MAG)

This supplementary material has been provided by the authors to give readers additional information about their work.

**eFigure.** Details of Patient Inclusion and Exclusion



**eTable 1.** Provider Volume and Baseline Patient Characteristics of Male Patients Undergoing Single-Arterial and Multiple-Arterial Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

Variable	Overall N (%)	Single arterial grafting N (%)	Multiple arterial grafting N (%)	SD
<b>Vessels diseased</b>				
• LM disease	13744 (28.5)	10948 (29.1)	2796 (26.6)	5.7
• Proximal LAD	26935 (55.9)	21188 (56.3)	5747 (54.6)	3.5
<b>2-vessel disease</b>				
• LAD and LCX	7166 (14.9)	5589 (14.9)	1577 (15.0)	0.4
• LAD and RCA	7743 (16.1)	6150 (16.3)	1593 (15.1)	3.3
• LCX and RCA	2803 (5.8)	2158 (5.7)	645 (6.1)	1.7
<b>3-vessel disease</b>	30443 (63.2)	23730 (63.1)	6713 (63.8)	1.4
<b>Previous AMI</b>				
• 0-23 Hours	407 (0.8)	340 (0.9)	67 (0.6)	3.1
• 1-7 Days	8821 (18.3)	7003 (18.6)	1818 (17.3)	3.5
• 8-14 Days	2039 (4.2)	1648 (4.4)	391 (3.7)	3.4
• 15-20 Days	382 (0.8)	323 (0.9)	59 (0.6)	3.6
• ≥21 Days	11314 (23.5)	9020 (24.0)	2294 (21.8)	5.2
<b>Left ventricular ejection fraction</b>				
• <20%	729 (1.5)	650 (1.7)	79 (0.8)	8.8
• 20-29%	3102 (6.4)	2649 (7.0)	453 (4.3)	11.9
• 30-39%	5585 (11.6)	4650 (12.4)	935 (8.9)	11.3
• 40-49%	9840 (20.4)	7809 (20.8)	2031 (19.3)	3.7
<b>CHF, current</b>	5043 (10.5)	4383 (11.6)	660 (6.3)	18.9
<b>CHF, past</b>	1399 (2.9)	1184 (3.1)	215 (2.0)	7.0
<b>Extensive aortic atherosclerosis</b>	2196 (4.6)	1796 (4.8)	400 (3.8)	4.8
<b>Hepatic failure</b>	8 (0.0)	7 (0.0)	1 (0.0)	0.8
<b>Renal dysfunction</b>				
• Serum creatinine 1.3-1.5	7218 (15.0)	5888 (15.6)	1330 (12.6)	8.7
• Serum creatinine 1.6-2.0	3209 (6.7)	2726 (7.2)	483 (4.6)	11.3
• Serum creatinine ≥2.1	1407 (2.9)	1231 (3.3)	176 (1.7)	10.3
<b>Renal dialysis</b>	1126 (2.3)	1048 (2.8)	78 (0.7)	15.6
<b>Previous PCI, current admission</b>	726 (1.5)	562 (1.5)	164 (1.6)	0.5
<b>Previous PCI, before current admission</b>	10205 (21.2)	7962 (21.2)	2243 (21.3)	0.4
<b>Surgeon average annual CABG Volume, &lt;75</b>	17477 (36.3)	14358 (38.2)	3119 (29.6)	18.1
<b>Surgeon average annual CABG Volume, 75-150</b>	22329 (46.4)	16075 (42.7)	6254 (59.4)	33.8

<b>Surgeon average annual CABG Volume, &gt;150</b>	8349 (17.3)	7194 (19.1)	1155 (11.0)	22.9
<b>Surgery priority</b>				
• <i>Urgent</i>	32023 (66.5)	25836 (68.7)	6187 (58.8)	20.7
• <i>Elective</i>	16132 (33.5)	11791 (31.3)	4341 (41.2)	20.7

AMI: acute myocardial infarction; CHF: congestive heart failure; LAD: left anterior descending artery; LCX: left circumflex artery; LM: left main; PCI: Percutaneous Coronary Intervention; RCA: right coronary artery; SD: standardized difference

**eTable 2.** Provider Volume and Baseline Patient Characteristics of Female Patients Undergoing Single-Arterial (SAG) and Multiple-Arterial (MAG) Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

Variable	Overall N (%)	Single arterial grafting N (%)	Multiple arterial grafting N (%)	SD
<b>Vessels diseased</b>				
• LM disease	4125 (27.1)	3568 (27.1)	557 (26.5)	1.4
• Proximal LAD	8400 (55.1)	7289 (55.4)	1111 (52.9)	5.2
<b>2-vessel disease</b>				
• LAD and LCX	2355 (15.4)	2035 (15.5)	320 (15.2)	0.7
• LAD and RCA	2710 (17.8)	2347 (17.9)	363 (17.3)	1.5
• LCX and RCA	920 (6.0)	776 (5.9)	144 (6.9)	3.9
<b>3-vessel disease</b>	9262 (60.7)	7988 (60.8)	1274 (60.6)	0.3
<b>Previous AMI</b>				
• 0-23 Hours	121 (0.8)	106 (0.8)	15 (0.7)	1.1
• 1-7 Days	3082 (20.2)	2708 (20.6)	374 (17.8)	7.1
• 8-14 Days	972 (6.4)	858 (6.5)	114 (5.4)	4.6
• 15-20 Days	182 (1.2)	160 (1.2)	22 (1.0)	1.6
• ≥21 Days	3064 (20.1)	2673 (20.3)	391 (18.6)	4.4
<b>Left ventricular ejection fraction</b>				
• <20%	137 (0.9)	126 (1.0)	11 (0.5)	5.1
• 20-29%	736 (4.8)	673 (5.1)	63 (3.0)	10.8
• 30-39%	1486 (9.7)	1327 (10.1)	159 (7.6)	8.9
• 40-49%	2670 (17.5)	2334 (17.8)	336 (16.0)	4.7
<b>CHF, current</b>	2381 (15.6)	2148 (16.3)	233 (11.1)	15.3
<b>CHF, past</b>	580 (3.8)	528 (4.0)	52 (2.5)	8.7
<b>Extensive aortic atherosclerosis</b>	1018 (6.7)	883 (6.7)	135 (6.4)	1.2
<b>Hepatic failure</b>	2 (0.0)	2 (0.0)	0 (0.0)	1.7
<b>Renal dysfunction</b>				
• Serum creatinine 1.3-1.5	1455 (9.5)	1292 (9.8)	163 (7.8)	7.3
• Serum creatinine 1.6-2.0	870 (5.7)	788 (6.0)	82 (3.9)	9.7
• Serum creatinine ≥2.1	477 (3.1)	423 (3.2)	54 (2.6)	3.9
<b>Renal dialysis</b>	433 (2.8)	409 (3.1)	24 (1.1)	13.7
<b>Previous PCI, current admission</b>	212 (1.4)	184 (1.4)	28 (1.3)	0.6
<b>Previous PCI, before current admission</b>	2862 (18.8)	2469 (18.8)	393 (18.7)	0.2
<b>Surgeon average annual CABG Volume, &lt;75</b>	5555 (36.4)	4976 (37.9)	579 (27.6)	22.1
<b>Surgeon average annual CABG Volume, 75-150</b>	7050 (46.2)	5716 (43.5)	1334 (63.5)	41.0

<b>Surgeon average annual CABG Volume, &gt;150</b>	2642 (17.3)	2454 (18.7)	188 (8.9)	28.5
<b>Surgery priority</b>				
• <i>Urgent</i>	10825 (71.0)	9451 (71.9)	1374 (65.4)	14.0
• <i>Elective</i>	4422 (29.0)	3695 (28.1)	727 (34.6)	14.0

AMI: acute myocardial infarction; CHF: congestive heart failure; LAD: left anterior descending artery; LCX: left circumflex artery; LM: left main; PCI: Percutaneous Coronary Intervention; RCA: right coronary artery; SD: standardized difference

**eTable 3.** Provider Volume and Baseline Characteristics of Propensity-Matched Male Patients Undergoing Single-Arterial and Multiple-Arterial Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

Variable	Overall N (%)	Single arterial grafting N (%)	Multiple arterial grafting N (%)	ASD
<b>Number of patients</b>	19024	9512	9512	-
<b>Age, years</b>	61.1 ( $\pm$ 9.5)	61.1 ( $\pm$ 9.5)	61.2 ( $\pm$ 9.4)	1.4
<b>Body surface area, m<sup>2</sup></b>	2.1 ( $\pm$ 0.2)	2.1 ( $\pm$ 0.3)	2.1 ( $\pm$ 0.2)	0.9
<b>Risk Score</b>	5.0 ( $\pm$ 2.5)	5.0 ( $\pm$ 2.5)	5.0 ( $\pm$ 2.5)	0.7
<b>High Risk (<math>\geq</math>9)</b>	1723 (9.1)	824 (8.7)	899 (9.5)	2.7
<b>Low Risk (&lt;9)</b>	17301 (90.9)	8688 (91.3)	8613 (90.5)	2.7
<b>Off-pump surgery</b>	3810 (20.0)	1915 (20.1)	1895 (19.9)	0.5
<b>Incomplete revascularization</b>	3466 (18.2)	1733 (18.2)	1733 (18.2)	0.0
<b>Number of conduits <math>\leq</math>3</b>	3791 (19.9)	1929 (20.3)	1862 (19.6)	1.8
<b>Age, <math>\geq</math> 70 years</b>	3566 (18.7)	1783 (18.7)	1783 (18.7)	0.0
<b>Vessels diseased</b>				
• <i>LM disease</i>	4578 (24.1)	2289 (24.1)	2289 (24.1)	0.0
• <i>Proximal LAD</i>	10466 (55.0)	5233 (55.0)	5233 (55.0)	0.0
<b>2-vessel disease</b>				
• <i>LAD and LCX</i>	2636 (13.9)	1318 (13.9)	1318 (13.9)	0.0
• <i>LAD and RCA</i>	2838 (14.9)	1408 (14.8)	1430 (15.0)	0.6
• <i>LCX and RCA</i>	1122 (5.9)	572 (6.0)	550 (5.8)	1.0
<b>3-vessel disease</b>	12428 (65.3)	6214 (65.3)	6214 (65.3)	0.0
<b>Previous AMI</b>				
• <i>0-23 Hours</i>	111 (0.6)	54 (0.6)	57 (0.6)	0.4
• <i>1-7 Days</i>	2901 (15.2)	1452 (15.3)	1449 (15.2)	0.1
• <i>8-14 Days</i>	666 (3.5)	296 (3.1)	370 (3.9)	4.2
• <i>15-20 Days</i>	98 (0.5)	47 (0.5)	51 (0.5)	0.6
• <i><math>\geq</math>21 Days</i>	4349 (22.9)	2216 (23.3)	2133 (22.4)	2.1
<b>Left ventricular ejection fraction</b>				
• <i>&lt;20%</i>	142 (0.7)	70 (0.7)	72 (0.8)	0.2
• <i>20-29%</i>	826 (4.3)	410 (4.3)	416 (4.4)	0.3
• <i>30-39%</i>	1658 (8.7)	833 (8.8)	825 (8.7)	0.3
• <i>40-49%</i>	3548 (18.7)	1766 (18.6)	1782 (18.7)	0.4
<b>Peripheral vascular disease</b>	1820 (9.6)	933 (9.8)	887 (9.3)	1.6
<b>Cerebrovascular disease</b>	2293 (12.1)	1147 (12.1)	1146 (12.0)	0.0
<b>CHF, current</b>	1161 (6.1)	582 (6.1)	579 (6.1)	0.1
<b>CHF, past</b>	418 (2.2)	221 (2.3)	197 (2.1)	1.7
<b>COPD</b>	2779 (14.6)	1443 (15.2)	1336 (14.0)	3.2
<b>Extensive aortic atherosclerosis</b>	697 (3.7)	352 (3.7)	345 (3.6)	0.4
<b>Diabetes</b>	5914 (31.1)	2999 (31.5)	2915 (30.6)	1.9

<b>Hepatic failure</b>	0 (0.0)	0 (0.0)	0 (0.0)	---
<b>Renal dysfunction</b>				
• <i>Serum creatinine 1.3-1.5</i>	2476 (13.0)	1223 (12.9)	1253 (13.2)	0.9
• <i>Serum creatinine 1.6-2.0</i>	614 (3.2)	297 (3.1)	317 (3.3)	1.2
• <i>Serum creatinine ≥2.1</i>	242 (1.3)	127 (1.3)	115 (1.2)	1.1
<b>Renal dialysis</b>	112 (0.6)	60 (0.6)	52 (0.5)	1.1
<b>Previous PCI, current admission</b>	287 (1.5)	142 (1.5)	145 (1.5)	0.3
<b>Previous PCI, before current admission</b>	4113 (21.6)	2098 (22.1)	2015 (21.2)	2.1
<b>Previous cardiac surgery</b>	304 (1.6)	150 (1.6)	154 (1.6)	0.3
<b>Surgeon average annual CABG Volume, &lt;75</b>	6038 (31.7)	3091 (32.5)	2947 (31.0)	3.3
<b>Surgeon average annual CABG Volume, 75-150</b>	10808 (56.8)	5358 (56.3)	5450 (57.3)	2.0
<b>Surgeon average annual CABG Volume, &gt;150</b>	2178 (11.4)	1063 (11.2)	1115 (11.7)	1.7
<b>Surgery priority</b>				
• <i>Urgent</i>	11241 (59.1)	5661 (59.5)	5580 (58.7)	1.7
• <i>Elective</i>	7783 (40.9)	3851 (40.5)	3932 (41.3)	1.7

AMI: acute myocardial infarction; ASD: adjusted standardized difference; CHF: congestive heart failure; COPD: chronic obstructive pulmonary disease; LAD: left anterior descending artery; LCX: left circumflex artery; LM: left main; PCI: Percutaneous Coronary Intervention; RCA: right coronary artery

**eTable 4.** Provider Volume and Baseline Characteristics of Propensity-Matched Female Patients Undergoing Single-Arterial (SAG) and Multiple-Arterial (MAG) Isolated-Coronary Artery Bypass Grafting (CABG) Surgery in New York State From 2005-2014

Variable	Overall N (%)	Single arterial grafting N (%)	Multiple arterial grafting N (%)	ASD
<b>Number of patients</b>	3720	1860	1860	-
<b>Age, years</b>	65.1 ( $\pm 10.3$ )	65.0 ( $\pm 10.3$ )	65.2 ( $\pm 10.3$ )	2.3
<b>Body surface area, m<sup>2</sup></b>	1.9 ( $\pm 0.2$ )	1.9 ( $\pm 0.2$ )	1.9 ( $\pm 0.2$ )	0.9
<b>Risk Score</b>	6.2 ( $\pm 2.6$ )	6.2 ( $\pm 2.5$ )	6.2 ( $\pm 2.7$ )	1.3
<b>High Risk (<math>\geq 9</math>)</b>	680 (18.3)	318 (17.1)	362 (19.5)	6.1
<b>Low Risk (&lt;9)</b>	3040 (81.7)	1542 (82.9)	1498 (80.5)	6.1
<b>Off-pump surgery</b>	778 (20.9)	369 (19.8)	409 (22.0)	5.3
<b>Incomplete revascularization</b>	782 (21.0)	391 (21.0)	391 (21.0)	0.0
<b>Number of conduits <math>\leq 3</math></b>	933 (25.1)	469 (25.2)	464 (24.9)	0.6
<b>Age, <math>\geq 70</math> years</b>	1278 (34.4)	639 (34.4)	639 (34.4)	0.0
<b>Vessels diseased</b>				
• <i>LM disease</i>	866 (23.3)	433 (23.3)	433 (23.3)	0.0
• <i>Proximal LAD</i>	1993 (53.6)	1009 (54.2)	984 (52.9)	2.7
<b>2-vessel disease</b>				
• <i>LAD and LCX</i>	506 (13.6)	253 (13.6)	253 (13.6)	0.0
• <i>LAD and RCA</i>	626 (16.8)	320 (17.2)	306 (16.5)	2.0
• <i>LCX and RCA</i>	226 (6.1)	106 (5.7)	120 (6.5)	3.2
<b>3-vessel disease</b>	2362 (63.5)	1181 (63.5)	1181 (63.5)	0.0
<b>Previous AMI</b>				
• <i>0-23 Hours</i>	16 (0.4)	7 (0.4)	9 (0.5)	1.6
• <i>1-7 Days</i>	540 (14.5)	271 (14.6)	269 (14.5)	0.3
• <i>8-14 Days</i>	196 (5.3)	92 (4.9)	104 (5.6)	2.9
• <i>15-20 Days</i>	36 (1.0)	17 (0.9)	19 (1.0)	1.1
• $\geq 21$ Days	733 (19.7)	376 (20.2)	357 (19.2)	2.6
<b>Left ventricular ejection fraction</b>				
• <20%	20 (0.5)	11 (0.6)	9 (0.5)	1.5
• 20-29%	101 (2.7)	50 (2.7)	51 (2.7)	0.3
• 30-39%	274 (7.4)	141 (7.6)	133 (7.2)	1.6
• 40-49%	559 (15.0)	274 (14.7)	285 (15.3)	1.7
<b>Peripheral vascular disease</b>	499 (13.4)	239 (12.8)	260 (14.0)	3.3
<b>Cerebrovascular disease</b>	719 (19.3)	354 (19.0)	365 (19.6)	1.5
<b>CHF, current</b>	375 (10.1)	182 (9.8)	193 (10.4)	2.0
<b>CHF, past</b>	93 (2.5)	47 (2.5)	46 (2.5)	0.3
<b>COPD</b>	754 (20.3)	383 (20.6)	371 (19.9)	1.6
<b>Extensive aortic atherosclerosis</b>	220 (5.9)	115 (6.2)	105 (5.6)	2.3
<b>Diabetes</b>	1614 (43.4)	800 (43.0)	814 (43.8)	1.5

<b>Hepatic failure</b>	0 (0.0)	0 (0.0)	0 (0.0)	---
<b>Renal dysfunction</b>				
• <i>Serum creatinine 1.3-1.5</i>	307 (8.3)	154 (8.3)	153 (8.2)	0.2
• <i>Serum creatinine 1.6-2.0</i>	108 (2.9)	54 (2.9)	54 (2.9)	0.0
• <i>Serum creatinine ≥2.1</i>	64 (1.7)	34 (1.8)	30 (1.6)	1.7
<b>Renal dialysis</b>	26 (0.7)	11 (0.6)	15 (0.8)	2.6
<b>Previous PCI, current admission</b>	46 (1.2)	23 (1.2)	23 (1.2)	0.0
<b>Previous PCI, before current admission</b>	718 (19.3)	365 (19.6)	353 (19.0)	1.6
<b>Previous cardiac surgery</b>	60 (1.6)	38 (2.0)	22 (1.2)	6.8
<b>Surgeon average annual CABG Volume, &lt;75</b>	1139 (30.6)	596 (32.0)	543 (29.2)	6.2
<b>Surgeon average annual CABG Volume, 75-150</b>	2240 (60.2)	1099 (59.1)	1141 (61.3)	4.6
<b>Surgeon average annual CABG Volume, &gt;150</b>	341 (9.2)	165 (8.9)	176 (9.5)	2.0
<b>Surgery priority</b>				
• <i>Urgent</i>	2384 (64.1)	1195 (64.2)	1189 (63.9)	0.7
• <i>Elective</i>	1336 (35.9)	665 (35.8)	671 (36.1)	0.7

AMI: acute myocardial infarction; ASD: adjusted standardized difference; CHF: congestive heart failure; COPD: chronic obstructive pulmonary disease; LAD: left anterior descending artery; LCX: left circumflex artery; LM: left main; PCI: Percutaneous Coronary Intervention; RCA: right coronary artery

**eTable 5.** Operative (In-Hospital/30 day) Mortality Rates of Female and Male Propensity Matched Patients

MAG: multiple arterial grafting; SAG: single arterial grafting

	Number of pairs	MAG (%)	SAG (%)	P-value (McNemar)
Female	2093	30 (1.43)	23 (1.10)	0.33
Male	10481	76 (0.73)	77 (0.73)	0.93

**eTable 6.** Seven-Year Mortality Rates of Propensity-Matched Risk-Stratified Female and Male Patients According to Different Low Risk Score Cut-Offs After Multiple- (MAG) vs Single-Arterial (SAG) Coronary Artery Bypass Grafting in New York State Between 2005-2014

Outcome	Multiple arterial grafting		Single arterial grafting		Adjusted hazard ratio (95% confidence interval)	P-value
	Patients at risk	Adverse event (KM-est)	Patients at risk	Adverse event (KM-est)		
<b>Female</b>						
Overall	999	362 (20.0)	972	386 (21.5)	0.92 (0.80-1.07)	0.29
Risk score (11 or less)	971	293 (17.5)	935	337 (20.0)	0.87 (0.74-1.02)	0.08
Risk score (10 or less)	940	252 (15.9)	900	301 (18.9)	0.83 (0.70-0.98)	0.03
Risk score (9 or less)	903	207 (13.9)	859	265 (17.7)	0.78 (0.65-0.94)	0.008
Risk score (8 or less)	869	183 (13.0)	834	233 (16.3)	0.80 (0.65-0.97)	0.02
Risk score (7 or less)	719	124 (11.0)	693	157 (13.5)	0.79 (0.62-1.00)	0.05
<b>Male</b>						
Overall	5408	1058 (11.9)	5301	1263 (14.2)	0.83 (0.77-0.90)	<0.001
Risk score (11 or less)	5354	919 (10.7)	5244	1125 (13.0)	0.83 (0.76-0.90)	<0.001
Risk score (10 or less)	5293	835 (9.9)	5183	1047 (12.4)	0.82 (0.74-0.89)	<0.001
Risk score (9 or less)	5244	796 (9.6)	5146	993 (11.9)	0.83 (0.75-0.91)	<0.001
Risk score (8 or less)	5081	681 (8.6)	4980	882 (11.1)	0.80 (0.73-0.89)	<0.001
Risk score (7 or less)	4562	504 (7.2)	4462	692 (9.8)	0.76 (0.67-0.85)	<0.001

KM est = Kaplan-Meier estimate

**eTable 7.** Fully Adjusted Cox Regression Models Evaluating the Effect of Multiple Arterial Grafting (MAG) on Death in the Overall Population

MAG: multiple arterial grafting

Variable	Hazard ratio MAG/SAG	P-value
MAG (hazard ratio MAG/SAG)	0.53 (0.46-0.61)	<0.001
Female (hazard ratio MAG/SAG)	1.44 (1.26-1.64)	<0.001
Risk score (hazard ratio MAG/SAG)	1.28 (1.27-1.29)	0.04
MAG-female sex interaction coefficient	1.13 (1.01-1.28)	<0.001
MAG-risk score interaction coefficient	1.06 (1.04-1.08)-	<0.001
Female sex-risk score interaction coefficient	0.96 (0.95-0.98)	<0.001

**eTable 8.** Landmark Analysis of Alive Female and Male Low- and High-Risk Patients at 30 Days

Patient group	Multiple arterial grafting		Single arterial grafting		Adjusted hazard ratio (95% confidence interval)	P-value
	Patients at risk	Survival (KM-est)	Patients at risk	Survival (KM-est)		
<b>Female</b>						
Low risk	862	171 (12.4)	828	223 (15.9)	0.78 (0.64-0.95)	0.01
High risk	126	163 (46.2)	136	141 (41.2)	1.06 (0.84-1.34)	0.61
<b>Male</b>						
Low risk	5058	633 (8.2)	4958	836 (10.7)	0.79 (0.71-0.88)	<0.001
High risk	322	348 (42.4)	313	352 (43.1)	0.94 (0.81-1.10)	0.46

KM est = Kaplan-Meier estimate

**eTable 9.** Sensitivity Analysis Using Gamma Testing

Comparing K-M survival curves for propensity-matched samples using Klein Moescherberger test

- Male: p<0.001
- Female: p=0.49

Comparing K-M survival free from major adverse cardiac and cerebrovascular event curves for propensity-matched samples using Klein Moescherberger test

- Male: p<0.001
- Female: p=0.94

7-year mortality

**Female**

Sensitivity Analysis for Klein and Moeschberger's Test

Gamma	P plus	P neg	P lower	P upper
-------	--------	-------	---------	---------

1.00	0.50000	0.50000	0.51864	0.51864
1.05	0.51220	0.48780	0.22530	0.93752
1.10	0.52381	0.47619	0.07957	1.00000
1.15	0.53488	0.46512	0.02323	1.00000
1.20	0.54545	0.45455	0.00572	1.00000
1.25	0.55556	0.44444	0.00121	1.00000

**Male**

Sensitivity Analysis for Klein and Moeschberger's Test

Gamma	P plus	P neg	P lower	P upper
-------	--------	-------	---------	---------

1.00	0.50000	0.50000	.000004877	0.00000
1.05	0.51220	0.48780	.000000020	0.00042
1.10	0.52381	0.47619	.000000000	0.01116
1.15	0.53488	0.46512	.000000000	0.11160
1.20	0.54545	0.45455	.000000000	0.49288
1.25	0.55556	0.44444	.000000000	1.00000

7-year major adverse cardiac and cerebrovascular events (MACCE)

**Female**

Sensitivity Analysis for Klein and Moeschberger's Test

Gamma	P plus	P neg	P lower	P upper
-------	--------	-------	---------	---------

1.00	0.50000	0.50000	0.97180	0.97180
1.05	0.51220	0.48780	0.46813	1.00000
1.10	0.52381	0.47619	0.16654	1.00000

1.15	0.53488	0.46512	0.04425	1.00000
1.20	0.54545	0.45455	0.00898	1.00000
1.25	0.55556	0.44444	0.00143	1.00000

**Male**

Sensitivity Analysis for Klein and Moeschberger's Test

Gamma	P plus	P neg	P lower	P upper
-------	--------	-------	---------	---------

1.00	0.50000	0.50000	.000054523	0.00005
1.05	0.51220	0.48780	.000000086	0.00657
1.10	0.52381	0.47619	.000000000	0.14374
1.15	0.53488	0.46512	.000000000	0.79244
1.20	0.54545	0.45455	.000000000	1.00000
1.25	0.55556	0.44444	.000000000	1.00000

**eTable 10.** Sensitivity Analysis Using E-Value

MACCE: Major adverse cardiac and cerebrovascular events

Outcome	E-value for point estimate	E-value for confidence interval
<b>Female</b>		
Mortality	1.09	1.00
MACCE	1.17	1.00
<b>Male</b>		
Mortality	1.61	1.44
MACCE	1.48	1.34

**eTable 11.** Fully Adjusted Cox Regression Models Comparing Mortality Between Female and Male Patients Receiving Single Arterial Grafts (SAG), and Female and Male Patients Receiving Multiple Arterial Grafts (MAG)

Patient group	Hazard ratio Female/Male	P-value
SAG	1.08 (1.04-1.13)	<0.001
MAG	1.22 (1.07-1.38)	<0.001