

1 Table S. Characteristic of patients included and excluded from the development of the nomogram ABL

Characteristics	Patients included (N=364)	Patients excluded (N = 378)	P Value
Demographics			
Age (median [IQR])	52.00 [41.00, 65.00]	52.00 [40.00, 64.00]	0.523
Gender male, n (%)	218 (59.9)	233 (61.6)	0.68
Ethnicity, n (%)			0.91
African American	28 (7.8)	26 (6.9)	
Asian	4 (1.1)	5 (1.3)	
Caucasian	274 (76.3)	293 (77.9)	
Hispanic	18 (5.0)	17 (4.5)	
Native American	3 (0.8)	6 (1.6)	
Other/Unknown	32 (8.9)	29 (7.7)	
Height (cm, median [IQR])	172.70 [163.00, 178.00]	172.00 [163.80, 178.00]	0.946
Weight (kg, median [IQR])	81.65 [69.00, 95.70]	86.19 [72.65, 100.15]	0.005
Clinical Factors			
ICU type, n (%)			0.267
Med-Surg ICU	252 (69.2)	253 (66.9)	
MICU	43 (11.8)	33 (8.7)	
SICU	28 (7.7)	37 (9.8)	
Other ICUs	41(11.3)	55(14.6)	
Comorbidities, n (%)			
CHF	22 (6.1)	21 (5.6)	0.912
COPD	39 (10.8)	35 (9.4)	0.605
Hypertension	163 (45.3)	177 (47.6)	0.582
DM	55 (15.3)	54 (14.5)	0.853
MI	12 (3.3)	16 (4.3)	0.624
RI	21 (5.8)	21 (5.6)	1
GCS (median [IQR])	15.00 [14.00, 15.00]	15.00 [14.00, 15.00]	0.513
In-hospital mortality	12 (3.3)	21 (5.6)	0.189

2 *Abbreviations:* IQR interquartile range, ICU intensive care unit, Med-Surg ICU medical-surgical intensive care unit, MICU medical
3 intensive care unit, SICU surgical intensive care unit, CHF chronic heart failure, COPD chronic obstructive pulmonary diseases,
4 DM diabetes mellitus, MI myocardial infarction, RI renal insufficiency, GCS Glasgow Coma Scale.