

Supporting Information for

NETosis Occurs Independently of Neutrophil Serine Proteases

Paulina Kasperkiewicz^{a,b}, Anne Hempel^a, Tomasz Janiszewski^b, Sonia Kołt^b, Scott J. Snipas^a, Marcin Drag^b, Guy S. Salvesen^{a,*}

Guy S. Salvesen

Email: gsalvesen@sbpdiscovery.org

This PDF file includes:

Legends for Movies S1 to S3c

Movie S1 Time course of NET formation in PMA-treated neutrophils co-incubated with a Cy5-labeled NE probe (red), and SYTOX Green™ to mark DNA.

Movie S2 Nuclear DNA is contained within the nucleus during pyroptosis induced by treatment of macrophages by LPS and ATP, and SYTOX Green™ to mark DNA

Movie S3a Z-stack of fresh neutrophils stimulated with PMA to induce NETosis BODIPY-labeled NE probe (green) and propidium iodide (red) to visualize DNA

Movie S3b Z-stack of fresh neutrophils stimulated with PMA to induce NETosis BODIPY-labeled CatG probe (green) and propidium iodide (red) to visualize DNA

Movie S3c Z-stack of fresh neutrophils stimulated with PMA to induce NETosis BODIPY-labeled PR3 probe (green) and propidium iodide (red) to visualize DNA