

RESEARCH SUMMARY

# Efficacy and Safety of mRNA-1273 SARS-CoV-2 Vaccine

L.R. Baden, et al. DOI: 10.1056/NEJMoa2035389

## CLINICAL PROBLEM

The Covid-19 pandemic continues and expands. Additional data regarding vaccines to prevent symptomatic severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection are needed. The mRNA-1273 vaccine is a lipid-encapsulated mRNA vaccine encoding the prefusion stabilized spike protein of SARS-CoV-2.

## CLINICAL TRIAL

A randomized, double-blind trial to evaluate the efficacy and safety of mRNA-1273.

30,420 participants ≥18 years old were assigned to receive either the vaccine or placebo in two intramuscular injections 28 days apart. Participants were followed for safety and the development of laboratory-confirmed, symptomatic Covid-19 over a median of 2 months after the second dose.

## RESULTS

### Safety:

Vaccine recipients had higher rates of local reactions (e.g., pain, erythema, swelling) and systemic reactions (e.g., headache, fatigue, myalgia) than placebo recipients. Most reactions were mild to moderate and resolved over 1–3 days.

### Efficacy:




The incidence of Covid-19 was lower among vaccine recipients than among placebo recipients as early as 14 days after the first dose. Protection in the vaccine group persisted for the period of follow-up.

## LIMITATIONS AND REMAINING QUESTIONS

Further study is required to understand the following:

- Safety and efficacy over a longer period of time, in a larger population, and in pregnant women and children.
- Whether the vaccine protects against asymptomatic infection and transmission to unvaccinated persons.
- How to care for those who miss the second vaccine dose.

Links: Full article | NEJM Quick Take | Editorial



	mRNA-1273 Vaccine N=14,550	Placebo N=14,598
Symptomatic Covid-19	11	185
Severe Covid-19	0	30

Vaccine efficacy of 94.1% (95% CI, 89.3–96.8%; P<0.001)

## CONCLUSIONS

Two doses of a SARS-CoV-2 mRNA-based vaccine were safe and provided 94% efficacy against symptomatic Covid-19 in persons 18 or older.