

Development of the Vision Impairment in Low Luminance (VILL) Questionnaire

Susanne G. Pondorfer, M.Sc. ^{1*}; Jan H. Terheyden, MD^{1*}; Helen Overhoff, M.Sc. ², Jana Stasch-Bouws, M.Sc.³, Frank G. Holz, MD¹; Robert P. Finger, MD, PhD¹

¹ Department of Ophthalmology, University Hospital Bonn, Bonn, Germany

² Forschungszentrum Jülich, Jülich, Germany

³ AMD-Netz e.V., Münster, Germany

SUPPLEMENT

Supplementary Table 1: Demographic characteristics of participants who were administered the VILL-68

	Overall	Early / intermediate AMD	Late AMD	<i>P</i>
<i>n</i>	127	74	53	
Age, Mean (SD)	76.1 (7.2)	76.0 (7.5)	76.3 (6.8)	0.809*
Gender, % (<i>n</i>)				0.846**
Female	69.3 (88)	70.3 (52)	67.9 (36)	
Male	30.7 (39)	29.7 (22)	32.1 (17)	
LogMAR visual acuity (better eye), Mean (SD)	0.1 (0.2)	0.1 (0.1)	0.2 (0.2)	0.022*
AMD stage, % (<i>n</i>)				
Early / intermediate AMD	58.3 (74)			
Late AMD	41.7 (53)			

* Unpaired t-test; ** Fisher's exact test