Supplemental material for:

Serbrouchni S, Malmartel A. Diagnostic agreement between telemedicine on social networks and teledermatology centers. *Ann Fam Med.* 2021;19(1):24-29.

Supplemental Table 1: complete list of diagnoses

	Expert committee	Twitter and MedPics	Teledermatology services
Viral rash	4 (6.7%)	4 (6.7%)	5 (8.3%)
Eczema	4 (6.7%)	5 (8.3%)	7 (11.7%)
Mycosis	4 (6.7%)	6 (10.0%)	6 (10.0%)
(mpetigo	2 (3.3%)	1 (1.7%)	1 (1.7%)
Photophytodermatosis	1 (1.7%)	2 (3.3%)	-
Non-tuberculous mycobacterium	1 (11,70)	1 (1.7%)	_
Purpura	5 (8.3%)	4 (6.7%)	4 (6.7%)
Syphilis	-	1 (1.7%)	-
Acrosyndrome	_	-	1 (1.7%)
Sarcoids	1 (1.7%)	1 (1.7%)	1 (1.7%)
Non conclusive diagnosis	4 (6.7%)	3 (5.0%)	3 (5.0%)
Granuloma annulare	-	1 (1.7%)	<i>5 (5.070)</i>
Orug allergy	1 (1.7%)	2 (3.3%)	1 (1.7%)
Scabies	1 (1.7%)	2 (3.3%)	- (1.770)
Herpes Labialis	3 (5.0%)	2 (3.3%)	3 (5.0%)
Hives	2 (3.3%)	2 (3.3%)	2 (3.3%)
Erythema migrans	3 (5.0%)	2 (3.3%)	2 (3.3%)
Psoriasis	1 (1.7%)	1 (1.7%)	2 (3.3%)
Vascularitis	1 (1.7%)	1 (1.7%)	1 (1.7%)
Prurigo	1 (1.770)	1 (1.7%)	1 (1.7%)
Lichen striatus	2 (3.3%)	2 (3.3%)	1 (1.7%)
Contact dermatitis	1 (1.7%)	1 (1.7%)	1 (1.770)
Morphea	1 (1.770)	1 (1.770)	1 (1.7%)
Skin cancer	2 (3.3%)	1 (1.7%)	1 (1.7%)
Palmoplantar keratoderma	2 (3.3%)	2 (3.3%)	2 (3.3%)
Lice and insect bites		2 (3.3%)	· · · · · · · · · · · · · · · · · · ·
Cushing syndrom	2 (3.3%)	1 (1.7%)	3 (5.0%)
e ·	-	1 (1.770)	1 (1.7%)
Polycythemia vera with erythrosis	- 1 (1 70/)	-	· /
Larva migrans	1 (1.7%)	1 (1 70/)	1 (1.7%)
Ulcerative tonsilitis	- 1 (1 7 0/)	1 (1.7%)	1 (1.7%)
Livedo Diamanta sania Kaban	1 (1.7%)	1 (1.7%)	1 (1.7%)
Pigmentogenic lichen	1 (1.7%)	1 (1.7%)	1 (1.7%)
Chalazion	1 (1.7%)	1 (1.7%)	- 2 (2 20/)
Molluscum contagiosum	1 (1.7%)	2 (3.3%)	2 (3.3%)
Herpes zoster	1 (1.7%)	1 (1.7%)	1 (1.7%)
Quincke's edema	1 (1.7%)	-	-
Flush	1 (1.7%)	-	-
Angina bullosa haemorrhagica	1 (1.7%)	-	1 (1 70/)
Linear IgA dermatosis	1 (1.7%)	-	1 (1.7%)
neutrophilic eccrine dermatosis	1 (1.7%)	-	-
Lymphangitis	1 (1.7%)	-	-
Chickenpox	1 (1.7%)	-	-
Folliculitis	1 (1.7%)	-	-
Biopsy required for a diagnosis	-	1 (1.7%)	3(5.0%)