nature research

Corresponding author(s): Dr. Carla Coffin

Last updated by author(s): Dr. Carla Coffin

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our <u>Editorial Policies</u> and the <u>Editorial Policy Checklist</u>.

Statistics

For a	statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.	
n/a	onfirmed	
	$rac{3}{3}$ The exact sample size (<i>n</i>) for each experimental group/condition, given as a discrete number and unit of measurement	
	🔾 A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly	
	C Only common tests should be described solely by name; describe more complex techniques in the Methods section.	
	A description of all covariates tested	
\boxtimes	A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons	
	A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals))
\boxtimes	For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted Give P values as exact values whenever suitable.	
\boxtimes	For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings	
\boxtimes	For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes	
\boxtimes	Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i>), indicating how they were calculated	
	Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.	

Software and code

Policy information about availability of computer code				
Data collection	NA			
Data analysis	Data was analysed using Graphpad Prism 6. Flow data was analyzed using FACS-DIVA and FlowJo softwares.			

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research guidelines for submitting code & software for further information.

Data

Policy information about availability of data

All manuscripts must include a <u>data availability statement</u>. This statement should provide the following information, where applicable: - Accession codes, unique identifiers, or web links for publicly available datasets

- A list of figures that have associated raw data
- A description of any restrictions on data availability

All the relevant information is in the manuscript in the form of tables, figures and supplementary material.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences

Behavioural & social sciences

Ecological, evolutionary & environmental sciences For a reference copy of the document with all sections, see <u>nature.com/documents/nr-reporting-summary-flat.pdf</u>

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	Unfortunately, there is limited data on HBV vaccine responses in NAFLD. The literature has shown that vaccination of 427 pre-adolescents, obesity was a significant predictor of poor immune response (P = 0.015) 4. In another study of 239 adult health care workers, older age (> 50 y) and diabetes was a poor predictor of non-response5. A study in cirrhotic patients (N=52) showed that cirrhosis, especially alcoholic chronic liver disease, was associated with lower antibody responses compared to the general population6. We initially performed a sample size calculation which stratified NAFLD patients according to normal BMI, overweight, obese and morbid obesity, as well as based on CAP score measurements, with intention to recruit ~200 patients. However, the study recruitment was more difficult than expected (>200 screen failure
Data exclusions	Data was not excluded
Replication	All cell culture measurements were carried out in triplicates.
Randomization	NA
Blinding	NA

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

- P	Л	et	۲ł	າດ	d	S

n/a	Involved in the study	n/a	Involved in the study
	Antibodies		ChIP-seq
	Eukaryotic cell lines		Flow cytometry
	Palaeontology and archaeology		MRI-based neuroimaging
	Animals and other organisms		
	Human research participants		
	Clinical data		
	Dual use research of concern		

Antibodies

Antibodies used	All antibodies have been described in supplementary table 1
Validation	Antibody concentrations were determined by titration especially for CCR7 and CD45RA. For the other highly expressed markers, data from the vendor (as specified on the technical data sheets) and core facility at University of Calgary was utilized

Eukaryotic cell lines

Policy information about <u>cell lines</u>	
Cell line source(s)	State the source of each cell line used.
Authentication	Describe the authentication procedures for each cell line used OR declare that none of the cell lines used were authenticated.
Mycoplasma contamination	Confirm that all cell lines tested negative for mycoplasma contamination OR describe the results of the testing for mycoplasma contamination OR declare that the cell lines were not tested for mycoplasma contamination.
Commonly misidentified lines (See <u>ICLAC</u> register)	Name any commonly misidentified cell lines used in the study and provide a rationale for their use.

Palaeontology and Archaeology

Specimen provenance	Provide provenance information for specimens and describe permits that were obtained for the work (including the name of the issuing authority, the date of issue, and any identifying information).
Specimen deposition	Indicate where the specimens have been deposited to permit free access by other researchers.
Dating methods	If new dates are provided, describe how they were obtained (e.g. collection, storage, sample pretreatment and measurement), where they were obtained (i.e. lab name), the calibration program and the protocol for quality assurance OR state that no new dates are provided.
Tick this box to confi	rm that the raw and calibrated dates are available in the paper or in Supplementary Information.
Ethics oversight	Identify the organization(s) that approved or provided guidance on the study protocol, OR state that no ethical approval or guidance was required and explain why not.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Animals and other organisms

Laboratory animals	Mice were purchased from Jackson Inc. (Bar Harbour, ME) and housed at the University of Calgary in a pathogen-free facility. All experiments were approved by the University of Calgary animal care committee in accordance with the Canadian Council for Animal care (AC16-0040). C57BL/6 male mice (6-8 weeks old, N=40;10/group)
Wild animals	Provide details on animals observed in or captured in the field; report species, sex and age where possible. Describe how animals were caught and transported and what happened to captive animals after the study (if killed, explain why and describe method; if released say where and when) OR state that the study did not involve wild animals.
Field-collected samples	For laboratory work with field-collected samples, describe all relevant parameters such as housing, maintenance, temperature, photoperiod and end-of-experiment protocol OR state that the study did not involve samples collected from the field.
Ethics oversight	University of Calgary animal care committee in accordance with the Canadian Council for Animal care (AC16-0040).

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Human research participants

Population characteristics	N=68 non-alcoholic fatty liver disease patients were recruited. Clinical characteristics are described in Table 1A
Recruitment	68 HBV naive NAFLD patients (18-60 years) in this study were recruited from specialized NAFLD clinics at 5 Canadian centres from August 2016 - August 2019. Exclusion criteria included subjects <18 and >60 years of age, pregnancy, human immunodeficiency virus+, HCV+, decompensated cirrhosis, and those with serological evidence of HBV exposure or immunization (HBsAg, anti-HBs) and total HBV core antibody (anti-HBc). All patients provided signed informed consent under an approved ethics protocol (Conjoint Health Research Ethics Board REB16-0274) according to the principles of Good Clinical Practice and the Declaration of Helsinki.
Ethics oversight	All patients provided signed informed consent under an approved ethics protocol (Conjoint Health Research Ethics Board REB16-0274) according to the principles of Good Clinical Practice and the Declaration of Helsinki.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Clinical data

Policy information about <u>clinical studies</u> All manuscripts should comply with the ICMJE guidelines for publication of clinical research and a completed CONSORT checklist must be included with all submissions. Clinical trial as sisteration. 1.7.1

Clinical trial registration	The study was registered on ClinicalTrials.gov (NCT02985450)			
Study protocol	https://clinicaltrials.gov/ct2/show/NCT02985450			
Data collection	Data were collected by clinical co-ordinators and clinicans at the recruiting centres at beaseline i.e. before vaccination and post vaccination. Clinical and demographic data were collected.			
Outcomes	Diagnosis of NAFLD was the primary outcome and determination of antibody titres in serum post 3 doses of vaccine as well as assessment of T-cell response in PBMC from these patients was the secondary outcome analyzed.			

Dual use research of concern

Policy information about <u>dual use research of concern</u>

Hazards

Could the accidental, deliberate or reckless misuse of agents or technologies generated in the work, or the application of information presented in the manuscript, pose a threat to:

No	Yes			
\ge		Public health		
\ge		National security		
\ge		Crops and/or livestock		
\ge		Ecosystems		
\ge		Any other significant area		
Experiments of concern				

Does the work involve any of these experiments of concern:

No	Yes
\boxtimes	Demonstrate how to render a vaccine ineffective
\boxtimes	Confer resistance to therapeutically useful antibiotics or antiviral agents
\boxtimes	Enhance the virulence of a pathogen or render a nonpathogen virulent
\boxtimes	Increase transmissibility of a pathogen
\boxtimes	Alter the host range of a pathogen
\boxtimes	Enable evasion of diagnostic/detection modalities
\boxtimes	Enable the weaponization of a biological agent or toxin
\boxtimes	Any other potentially harmful combination of experiments and agents

ChIP-seq

Data deposition

Confirm that both raw and final processed data have been deposited in a public database such as GEO.

Confirm that you have deposited or provided access to graph files (e.g. BED files) for the called peaks.

Data access links May remain private before publication.	For "Initial submission" or "Revised version" documents, provide reviewer access links. For your "Final submission" document, provide a link to the deposited data.
Files in database submission	Provide a list of all files available in the database submission.
Genome browser session (e.g. <u>UCSC</u>)	Provide a link to an anonymized genome browser session for "Initial submission" and "Revised version" documents only, to enable peer review. Write "no longer applicable" for "Final submission" documents.

Methodology

Replicates	Describe the experimental replicates, specifying number, type and replicate agreement.
Sequencing depth	Describe the sequencing depth for each experiment, providing the total number of reads, uniquely mapped reads, length of reads and whether they were paired- or single-end.
Antibodies	Describe the antibodies used for the ChIP-seq experiments; as applicable, provide supplier name, catalog number, clone name, and lot number.
Peak calling parameters	Specify the command line program and parameters used for read mapping and peak calling, including the ChIP, control and index files used.
Data quality	Describe the methods used to ensure data quality in full detail, including how many peaks are at FDR 5% and above 5-fold enrichment.
Software	Describe the software used to collect and analyze the ChIP-seq data. For custom code that has been deposited into a community repository, provide accession details.

Flow Cytometry

Plots

Confirm that:

The axis labels state the marker and fluorochrome used (e.g. CD4-FITC).

The axis scales are clearly visible. Include numbers along axes only for bottom left plot of group (a 'group' is an analysis of identical markers).

All plots are contour plots with outliers or pseudocolor plots.

A numerical value for number of cells or percentage (with statistics) is provided.

Methodology

Sample preparation	PBMC were separated by density gradient centrifugation from ~20 mL of heparinized blood, and ~107 cells/vial were cryopreserved. PBMC were only isolated at the coordinating centre at baseline (i.e. before the vaccination) and post vaccination for assessment of cellular response.
Instrument	BD FACS Canto II, Toronto, ON, Canada
Software	Data was analyzed using FACS DIVA and Flowjo v11 (Treestar Inc, San Carlos, CA).
Cell population abundance	Lymphocytes were gated based on FSC-SSC profiles, followed by identification of singlets, viable cells, CD3+CD56- cells, followed by identification of CD4+ and CD8+T cells and lastly identifying Memory T cells based on CCR7 and CD45RA. CD4+ T cells were also drilled down to identify proliferation using CFSE.
Gating strategy	Figure 3A has a representative gating strategy of immunophenotyping of memory T cells and proliferation of CD4+ T cells.
Tick this box to confirm th	at a figure exemplifying the gating strategy is provided in the Supplementary Information.

Magnetic resonance imaging

Experimental design

Experimental design	
Design type	Indicate task or resting state; event-related or block design.
Design specifications	Specify the number of blocks, trials or experimental units per session and/or subject, and specify the length of each trial or block (if trials are blocked) and interval between trials.
Behavioral performance measure	es State number and/or type of variables recorded (e.g. correct button press, response time) and what statistics were used to establish that the subjects were performing the task as expected (e.g. mean, range, and/or standard deviation across subjects).
Acquisition	
Imaging type(s)	Specify: functional, structural, diffusion, perfusion.
Field strength	Specify in Tesla
Sequence & imaging parameters	Specify the pulse sequence type (gradient echo, spin echo, etc.), imaging type (EPI, spiral, etc.), field of view, matrix size, slice thickness, orientation and TE/TR/flip angle.
Area of acquisition	State whether a whole brain scan was used OR define the area of acquisition, describing how the region was determined.
Diffusion MRI Used	Not used
Preprocessing	
Preprocessing software	Provide detail on software version and revision number and on specific parameters (model/functions, brain extraction, segmentation, smoothing kernel size, etc.).
Normalization	If data were normalized/standardized, describe the approach(es): specify linear or non-linear and define image types used for transformation OR indicate that data were not normalized and explain rationale for lack of normalization.
Normalization template	Describe the template used for normalization/transformation, specifying subject space or group standardized space (e.g. original Talairach, MNI305, ICBM152) OR indicate that the data were not normalized.

physiological signals (heart rate, respiration).

Describe your procedure(s) for artifact and structured noise removal, specifying motion parameters, tissue signals and

Noise and artifact removal

Define your software and/or method and criteria for volume censoring, and state the extent of such censoring.

Statistical modeling & inference

Model type and settings	Specify type (mass univariate, multivariate, RSA, predictive, etc.) and describe essential details of the model at the first and second levels (e.g. fixed, random or mixed effects; drift or auto-correlation).			
Effect(s) tested	Define precise effect in terms of the task or stimulus conditions instead of psychological concepts and indicate whether ANOVA or factorial designs were used.			
Specify type of analysis: Whole brain ROI-based Both				
Statistic type for inference (See <u>Eklund et al. 2016</u>)	specify voxel wise of elaster wise and report an relevant parameters for elaster wise methods.			
Correction	Describe the type of correction and how it is obtained for multiple comparisons (e.g. FWE, FDR, permutation or Monte Carlo).			

Models & analysis

- n/a Involved in the study
- Functional and/or effective connectivity
- Graph analysis
- Multivariate modeling or predictive analysis