

**REPEATED ELECTROMAGNETIC FIELD STIMULATION LOWERS AMYLOID-
BETA PEPTIDE LEVELS IN PRIMARY HUMAN NEURON-RICH BRAIN TISSUE
CULTURES**

Felipe P. Perez,² Bryan Maloney,¹ Nipun Chopra,¹ Jorge J. Morisaki,³ and Debomoy K. Lahiri¹

¹Department of Psychiatry, Institute of Psychiatric Research, Neuroscience Research Center, Indiana University School of Medicine, Indianapolis, IN, USA.

²Department of Medicine, Division of General Internal Medicine and Geriatrics, Aging for Center Research, Indiana University School of Medicine, Indianapolis, IN, USA.

³Department of Bioengineering, University of Illinois at Chicago, Chicago, IL, USA.

Corresponding author

Dr. D. K Lahiri

Neuroscience Research Center

320 W. 15th St., Indianapolis IN 46201, USA

Tel: (317) 274-2706

E-mails: dlahiri@iupui.edu

REMF reduces A β




Fig. S1. Total sAPP was measured by semiquantitative western blotting of CM from PHB-DIV28 exposed to REMFS at 64 MHz with SAR of 0.9 W/Kg for 4 or 8 days, as described in the main text. Un-edited scan of X-ray film with superimposed protein MW standard (BioRad 1610374) scanned from Ponceau-S stain of blot before antibody probing.