

Supplementary Digital Content 1. Participant-reported Pain Diagnoses within the “Other” category

Diagnoses, <i>n</i> (%)	Initial Survey (Study 1) <i>n</i> = 530	Follow-up Survey (Study 2) <i>n</i> = 278	In-person Interviews (Study 3) <i>n</i> = 28
Abdominal Pain	6 (1.1%)	0 (0.0%)	0 (0.0%)
Ankylosing Spondylitis	3 (0.6%)	1 (0.3%)	0 (0.0%)
Arthritis	8 (1.5%)	7 (2.5%)	0 (0.0%)
Arthromyalgia	0 (0.0%)	0 (0.0%)	1 (3.6%)
Autoimmune Disease	6 (1.1%)	3 (1.1%)	0 (0.0%)
Avascular Necrosis	1 (0.2%)	0 (0.0%)	0 (0.0%)
Back Pain	18 (3.4%)	4 (1.4%)	3 (10.7%)
Bursitis	2 (0.4%)	0 (0.0%)	0 (0.0%)
Cancer Pain	0 (0.0%)	0 (0.0%)	1 (3.6%)
Central Pain Syndrome	0 (0.0%)	0 (0.0%)	1 (3.6%)
Cervical Stenosis	1 (0.2%)	0 (0.0%)	0 (0.0%)
Chronic Fatigue Syndrome	4 (0.8%)	2 (0.7%)	0 (0.0%)
Crohn’s Disease	1 (0.2%)	0 (0.0%)	0 (0.0%)
Complex Regional Pain Syndrome (CRPS)	0 (0.0%)	1 (0.3%)	0 (0.0%)
Degenerative Disc Disease	6 (1.1%)	4 (1.4%)	2 (7.2%)
Dystonia	3 (0.6%)	2 (0.7%)	0 (0.0%)
Ehlers Danlos Syndrome	12 (2.3%)	7 (2.5%)	1 (3.6%)
Failed Back Surgery Syndrome	11 (2.1%)	4 (1.4%)	1 (3.6%)
Fibromyalgia	2 (0.4%)	0 (0.0%)	0 (0.0%)
Headache	8 (1.5%)	13 (4.6%)	0 (0.0%)
Herniated Disc	3 (0.6%)	0 (0.0%)	0 (0.0%)
Inflammatory Pain	1 (0.2%)	4 (1.4%)	0 (0.0%)
Injury, unspecified	18 (3.4%)	5 (1.8%)	1 (3.6%)
Interstitial Cystitis	1 (0.2%)	0 (0.0%)	0 (0.0%)
Kidney Pain	2 (0.4%)	0 (0.0%)	0 (0.0%)
Lyme Disease	1 (0.2%)	0 (0.0%)	0 (0.0%)
Lymphedema	1 (0.2%)	0 (0.0%)	0 (0.0%)
Marfan’s Causing Arthritis	1 (0.2%)	0 (0.0%)	0 (0.0%)
Meralgia Paresthetica	2 (0.4%)	1 (0.3%)	0 (0.0%)
Migraine	5 (0.9%)	0 (0.0%)	0 (0.0%)
Musculoskeletal Pain	24 (4.5%)	22 (7.9%)	0 (0.0%)

Myasthenia Gravis	1 (0.2%)	1 (0.3%)	0 (0.0%)
Myofascial Pain	3 (0.6%)	1 (0.3%)	0 (0.0%)
Neck Pain	4 (0.8%)	2 (0.7%)	0 (0.0%)
Neuropathic Pain	20 (3.8%)	8 (2.9%)	1 (3.6%)
Occipital Neuralgia	3 (0.6%)	3 (1.1%)	0 (0.0%)
Osteoarthritis	2 (0.4%)	0 (0.0%)	0 (0.0%)
Osteoporosis	1 (0.2%)	0 (0.0%)	0 (0.0%)
Pelvic Pain /Vulvodynia	5 (0.9%)	2 (0.7%)	0 (0.0%)
Phantom Limb Pain	2 (0.4%)	0 (0.0%)	0 (0.0%)
Post-polio Syndrome	1 (0.2%)	1 (0.3%)	0 (0.0%)
Postsurgical Pain	13 (2.5%)	10 (3.6%)	1 (3.6%)
Postherpetic Neuralgia (Shingles)	2 (0.4%)	0 (0.0%)	0 (0.0%)
Postural Orthostatic Tachycardia Syndrome (POTS)	1 (0.2%)	0 (0.0%)	0 (0.0%)
Psychosomatic	2 (0.4%)	0 (0.0%)	0 (0.0%)
Radiculopathy	9 (1.7%)	6 (2.1%)	0 (0.0%)
Sciatica	1 (0.2%)	0 (0.0%)	0 (0.0%)
Shoulder Pain	1 (0.2%)	0 (0.0%)	0 (0.0%)
SI Joint Dysfunction	1 (0.2%)	0 (0.0%)	0 (0.0%)
Sjogren's Syndrome	0 (0.0%)	1 (0.3%)	0 (0.0%)
Spinal Stenosis	0 (0.0%)	2 (0.7%)	0 (0.0%)
Tarlov Cyst	3 (0.6%)	1 (0.3%)	0 (0.0%)
Tethered Cord	1 (0.2%)	0 (0.0%)	0 (0.0%)
Thoracic Outlet Syndrome	0 (0.0%)	1 (0.3%)	0 (0.0%)
Trigeminal Neuralgia	1 (0.2%)	0 (0.0%)	0 (0.0%)
Tumor Pain	2 (0.4%)	0 (0.0%)	0 (0.0%)
Vascular Pain	3 (0.6%)	0 (0.0%)	0 (0.0%)
Vulvodynia	1 (0.2%)	0 (0.0%)	0 (0.0%)

*Percentages are based on the total cohort sample size