

**Early Chronic Kidney Disease Care Program delays kidney function  
deterioration in patients with Stage I–IIIa chronic kidney disease: An  
observational cohort study in Taiwan**

Shu-Fen Niu, Chung-Kuan Wu, Nai-Chen Chuang, Ya-Bei Yang, Tzu-Hao Chang

**Supplementary material**

**Table S1.** International Classification of Diseases, ninth Revision, Clinical

Modification codes used to identify comorbid conditions in this study.

**Table S2.** Distribution of eGFR amongst cases and controls during the follow-up period.

**Table S1.** International Classification of Diseases, ninth Revision, Clinical

Modification codes used to identify comorbid conditions in this study.

Co-morbid diseases	Corresponding ICD-9-CM codes
Hypertension	401.x–405.x
Diabetes mellitus	250.x
Gout	274.x
Hyperlipidemia	272.x
Heart disease	410.x–414.x; 420.x–429.x
Cerebrovascular disease	430.x–438.x

**Table S2.** Distribution of eGFR amongst cases and controls during the follow-up

period.

	eGFR				
	Control (n = 2076)		Case (n = 1038)		p-value
	N	Mean ± SD	N	Mean ± SD	
Follow-up time					
Baseline	2076	62.2 ± 13.3	1038	62.2 ± 12.9	0.8842
1 year	479	67.0 ± 26.0	398	63.3 ± 17.9	<b>0.0122</b>
2 years	385	65.4 ± 24.5	239	65.5 ± 19.6	0.9406
3 years	301	63.0 ± 24.1	143	66.5 ± 19.3	0.0970
5 years	181	56.7 ± 20.9	17	59.1 ± 13.9	0.6338