



Article

Identification of Molecular Basis for Objective Discrimination of Breast Cancer Cells (MCF-7) from Normal Human Mammary Epithelial Cells by Raman Microspectroscopy and Multivariate Curve Resolution Analysis

Keita Iwasaki ¹, Asuka Araki ², Murali Krishna C ³, Riruke Maruyama ², Tatsuyuki Yamamoto ^{4,5,*} and Hemanth Noothalapati ^{5,6,*}

¹ The United Graduate School of Agricultural Sciences, Tottori University, Tottori 680-8550, Japan; d17a3003@matsu.shimane-u.ac.jp

² Department of Organ Pathology, Faculty of Medicine, Shimane University, Izumo 693-8501, Japan; asuka@med.shimane-u.ac.jp (A.A.); hm5995@med.shimane-u.ac.jp (R.M.)

³ Advanced Centre for Treatment, Research and Education in Cancer, Tata Memorial Centre, 410-210 Navi Mumbai, India; pittu1043@gmail.com

⁴ Faculty of Life and Environmental Science, Shimane University, Matsue 690-8504, Japan

⁵ Raman Project Center for Medical and Biological Applications, Shimane University, Matsue 690-8504, Japan

⁶ Research Administration Office, Shimane University, Matsue 690-8504, Japan

* Correspondence: tyamamot@life.shimane-u.ac.jp (T.Y.); nvhnag@life.shimane-u.ac.jp (H.N.)

Supplementary Information




Figure S1. Preliminary MCR analysis of whole fingerprint region (1800–370 cm^{-1}). No significant bands were observed below 750 cm^{-1} .