

THE LANCET

Supplementary appendix

This appendix formed part of the original submission. We post it as supplied by the authors.

Supplement to: Crovetto F, Crispi F, Llurba E, Figueras F, Gómez-Roig MD, Gratacós E. Seroprevalence and presentation of SARS-CoV-2 in pregnancy. *Lancet* 2020; published online August 6. [http://dx.doi.org/10.1016/S0140-6736\(20\)31714-1](http://dx.doi.org/10.1016/S0140-6736(20)31714-1).

1 **Appendix**


Table. Baseline characteristics and clinical features of pregnant women positive for SARS-CoV-2 infection (N=125) in the first vs. third trimester.

Characteristics	First Trimester	Third Trimester	p
	N=54	N=71	
Age (years)	33.7 (5.1)	31.8 (6.1)	0.062
White ethnicity	31 (57)	43 (61)	0.697
Body mass index (Kg/m ²)	24.1 (6.2)	23.9 (4.3)	0.209
Smoking status	4 (7)	6 (9)	0.543
Medical history			
Chronic hypertension	1 (2)	2 (3)	0.602
Diabetes mellitus	1 (2)	0 (0)	0.432
Autoimmune disease	2 (4)	0 (0)	0.185
Respiratory disease	2 (4)	3 (4)	0.297
Nulliparous	34 (63)	36 (51)	0.095
Multiple pregnancy	2 (4)	1 (1)	0.398
Symptoms compatible with COVID-19 within last 2months			
None	38 (70)	37 (52)	0.012
Fever	7 (13)	16 (23)	0.152
Dry cough	5 (9)	17 (24)	0.034
Dyspnea	0 (0)	8 (11)	0.010
Tiredness	5 (9)	4 (6)	0.307
Sore throat	6 (11)	3 (4)	0.116
Rash on skin, or discoloration of fingers/toes	0 (0)	0 (0)	NA
Diarrhea	4 (7)	5 (7)	0.575
Loss of taste or smell	7 (13)	16 (23)	0.152
At least 3 symptoms or anosmia	9 (17)	22 (31)	0.065
Fever, cough and dyspnea	0 (0)	8 (11)	0.010
Hospital admission for COVID-19 symptoms	0 (0)	7 (10)	0.019

SARS-CoV-2: severe acute respiratory syndrome coronavirus 2

Data are expressed as n (%) or mean (SD)

3 **Figure.** Clinical spectrum of SARS-CoV-2 infection in pregnant women in the first and third trimester.
4



5
6