miR-361-3p Regulates Liver Tumor-initiating Cells Expansion and

Chemo-resistance

Shuping Qu^{1,#}, Xiaobing Zhang^{1,#}, Yue Wu^{1,#}, HengYu Li², Jian Zhai^{1,*}, Dong Wu^{1,*}

¹Department of Hepatic Surgery, Third Affiliated Hospital of Second Military Medical University, Shanghai, 200438, China.

²Department of General surgery, First Affiliated Hospital of Second Military Medical University, Shanghai, 200433, China.

[#]These authors contributed equally to this work.


Correspondence author address: Department of Hepatic Surgery, Third Affiliated Hospital of Second Military Medical University, Shanghai, 200438, China, E-mail: jianzhai1979@126.com (Jian Zhai); <u>WuDongEHBH@126.com</u> (Dong Wu).

Running title: miR-361-3p promotes liver T-ICs expansion.

Funding

This work is supported by the National Natural Science Foundation of China (81672718).

Supplementary Figure 1



Supplementary Figure 1:

A. The expression of SOX1 in CD24⁺ and CD24⁻ HCC cells was assessed by real-time PCR assay.

B. The expression of SOX1 in CD133⁺ and CD133⁻ HCC cells was detected by real-time PCR assay.

C. The expression of SOX1 in EpCAM⁺ and EpCAM⁻ HCC cells was checked by real-time PCR assay.

D. The expression of SOX1 in hepatoma spheroids was examined by real-time PCR assay.

E. The expression of SOX1 in CD24⁺ and CD24⁻ primary HCC cells was assessed by

real-time PCR assay.

F. The expression of SOX1 in CD133⁺ and CD133⁻ primary HCC cells was detected by real-time PCR assay.

G. The expression of SOX1 in EpCAM⁺ and EpCAM⁻ primary HCC cells was checked by real-time PCR assay.

H. The expression of SOX1 in primary hepatoma spheroids was examined by real-time PCR assay.

Supplementary Tables

Characteristics		Total n-70
Characteristics		Total n=70
Age(year)	≤ 50	44
	>50	26
Gender	Male	65
	Female	5
HBsAg	Positive	57
	Negative	13
AFP(µg/L)	≤400	51
	>400	19
T . ()	≤5	26
Tumor size(cm)	>5	44
Tumor number	Single	43
	Multiple	27
	Yes	12
Portal vein tumor thrombus	No	58
Pathological satellite	Yes	28
	No	42
BCLC stage	А	29
	B or C	41
	I-II	30
TNM	I-II	40

Supplementary Table 1. Clinicopathological features of 70 HCC patients in Cohort 1

HBsAg: hepatitis B virus surface antigen; AFP: α-fetoprotein; TNM: Tumor-Nodes-Metastasis; BCLC: Barcelona Clinic Liver Cancer Staging.

Characteristics		Total n=68
Age(year)	≤50	40
	>50	28
Gender	Male	61
	Female	7
	Positive	57
HBsAg	Negative	11
$\Delta ED(u a/L)$	≤400	50
$AFP(\mu g/L)$	>400	18
Tumor size(am)	≤5	19
Tumor size(cm)	>5	49
Tumor number	Single	38
	Multiple	30
Dortal yoin tymor thrombus	Yes	10
Portal vein tumor thrombus	No	58
Pathological satellite	Yes	25
	No	43
	А	26
BCLC stage	B or C	42
TNM	I-II	29
I INIVI	I-II	39

Supplementary Table 2. Clinicopathologic Features of 68 HCC Specimens in

Cohort 2

HBsAg: hepatitis B virus surface antigen; AFP: α-fetoprotein; TNM: Tumor-Nodes-Metastasis; BCLC: Barcelona Clinic Liver Cancer Staging.

Gene	Forward primer	Reverse primer (5'-3')	
miR-361-3p	Forward	GCCGCTCCCCAGGTGTGATT	
	Reverse	GTGCAGGGTCCGAGGT	
U6	Forward	CTCGCTTCGGCAGCACATA	
	Reverse	AACGATTCACGAATTTGCGT	
β-actin	Forward	GGCCCAGAATGCAGTTCGCCTT	
	Reverse	AATGGCACCCTGCTCACGCA	
SOX2	Forward	TGGAGAAGGAATGGTCCACTTC	
	Reverse	GGATAAGTACACGCTGCCCG	
OCT4	Forward	ATGTGCGCGTAACTGTCCAT	
	Reverse	CTGCAGTGTGGGGTTTCGGGGCA	
SOX1	Forward	AATACTGGAGACGAACGCCG	
	Reverse	CCCTCGAGCAAAGAAAACGC	
	Forward	CCCTCCACTCGGAAGGACTA	
c-Myc	Reverse	GCTGGTGCATTTTCGGTTGT	

Supplementary Table 3. Primer List.

Supplementary Table 4. Antibody List.

Antigens	Manufacturer	Application
SOX1	Proteintech Group, China	1:1000 for WB
PARP	Cell Signaling Technology, Beverly, MA	1:1000 for WB
β-actin	Santa Cruz Biotechnology, CA	1:5000 for WB
GAPDH	Santa Cruz Biotechnology, CA	1:5000 for WB