

**Table S2:** Non-treatment naïve/pubertal patient characteristics at baseline, with or without Laron syndrome (Enrolled population)

	Non-NPP-Laron syndrome (N = 17)		Non-NPP-non-Laron syndrome (N = 87)		P values <sup>a</sup>
	N <sup>b</sup>		N <sup>b</sup>		
<b>Female, n (%)</b> 95% CI	17	7 (41.2) 21.6; 64.0	87	28 (32.2) 23.3; 42.6	0.473 <sup>c</sup>
<b>Age at first injection, years</b> Mean (SD [95% CI]) Median (Q1; Q3)	17	12.78 (3.73 [10.86; 14.70]) 12.16 (10.72; 15.29)	87	11.43 (3.58 [10.67; 12.20]) 12.25 (8.74; 13.93)	0.272 <sup>w</sup>
<b>Height, SDS</b> Mean (SD [95% CI]) Median (Q1; Q3)	15	-4.63 (1.51 [-5.47; -3.79]) -4.93 (-5.89; -3.30)	77	-3.61 (1.20 [-3.88; -3.34]) -3.25 (-4.33; -2.84)	0.011 <sup>w</sup>
<b>Height velocity, cm/year</b> Mean (SD [95% CI]) Median (Q1; Q3)	12	4.43 (1.23 [3.65; 5.21]) 4.57 (3.69; 5.18)	52	4.70 (1.84 [4.19; 5.21]) 4.53 (3.70; 5.60)	0.884 <sup>w</sup>
<b>Weight, SDS</b> Mean (SD [95% CI]) Median (Q1; Q3)	15	-2.34 (1.28 [-3.05; -1.63]) -2.42 (-3.37; -1.17)	79	-3.28 (1.54 [-3.62; -2.93]) -2.92 (-4.10; -2.34)	0.036 <sup>w</sup>
<b>BMI, SDS</b>	15		71		<0.001 <sup>w</sup>

Mean (SD [95% CI])		0.86 (1.56 [-0.00; 1.73])		-0.97 (1.38 [-1.30; -0.65])	
Median (Q1; Q3)		0.81 (-0.45; 1.85)		-0.93 (-1.57; -0.27)	
<b>Mother's height, cm</b>	16		76		-
Mean (SD [95% CI])		158.1 (10.2 [152.7; 163.5])		157.5 (9.2 [155.4; 159.6])	
Median (Q1; Q3)		158.1 (148.7; 167.5)		158.0 (151.5; 163.6)	
<b>Father's height, cm</b>	17		76		-
Mean (SD [95% CI])		173.2 (5.8 [170.2; 176.2])		171.4 (8.2 [169.6; 173.3])	
Median (Q1; Q3)		172.0 (170.0; 178.0)		171.5 (165.3; 177.5)	
<b>IGF-1, ng/mL</b>	13		78		0.153 <sup>w</sup>
Mean (SD [95% CI])		261.41 (205.25 [137.38; 385.44])		148.74 (144.66 [116.12; 181.35])	
Median (Q1; Q3)		246.00 (62.00; 462.00)		105.50 (60.00; 171.10)	
<b>Peak stimulated GH level, ng/mL</b>	9		52		<0.001 <sup>w</sup>
Mean (SD [95% CI])		62.77 (37.09 [34.25; 91.28])		23.95 (50.31 [9.94; 37.96])	
Median (Q1; Q3)		55.00 (40.00; 72.90)		15.10 (7.45; 23.25)	
<b>Primary diagnosis: SPIGFD<sup>†</sup>, n (%)</b>	17	17 (100)	87	63 (72.4)	-
95% CI		81.6; 100		62.2; 80.7	
<b>History of hypoglycaemia, n (%)</b>	17	5 (29.4)	87	5 (5.7)	0.010 <sup>f</sup>
95% CI		13.3; 53.1		2.5; 12.8	

Data are median Q1; Q3 unless stated otherwise. <sup>†</sup>including Laron syndrome, as reported by the investigator. <sup>a</sup>Non-NPP-Laron syndrome versus Non-NPP-non-Laron syndrome. <sup>b</sup>Number of patients with available data; (c) Chi-square test; (f) Fisher's test; (w) Wilcoxon test. BMI, body mass index; GH, growth

hormone; IGF-1; insulin-like growth factor-1; NPP, treatment-naïve and prepubertal; SD, standard deviation; SDS, standard deviation score; SPIGFD, severe primary IGF-1 deficiency.