

Appendix 3 (as supplied by the authors): Supplemental tables

Supplemental Table S1: Quality of serious illness conversations based on chart review using a validated codebook

Domain	Intervention (n=55)	Control (n=28)	P-value
Life-sustaining treatments, total (/2)	2 (1,2)	1 (0,2)	0.03
Prognosis and illness understanding, total (/4)	3 (2,3)	1 (0,2)	<0.001
Values and goals, total (/7)	4 (2,6)	1 (0,1)	<0.001
EOL care planning, total (/4)	0.5 (0,2)	2 (0,2)	0.48
Overall (/17)	9 (7,12)	4 (3,6)	<0.001

All values reported as median (interquartile range) mean (SD)

Supplemental Table S2: Quality of Serious Illness Conversations: presence of individual codebook items

Domain		Intervention (n=55)	Control (n=28)	P-value
Life-sustaining treatments	Code status	42 (76)	20 (71)	0.62
	Life-sustaining treatments, etc	37 (67)	9 (32)	0.002
Prognosis and illness understanding	Illness understanding	45 (82)	8 (29)	<0.001
	Information preferences	18 (33)	0 (0)	0.001
	Prognosis, unspecified	47 (85)	19 (68)	0.06
	Prognosis communication about time, etc	34 (62)	8 (29)	0.004
Values and goals	Goals of care, unspecified	46 (84)	15 (54)	0.003
	Goals, priorities, "is important to"	38 (69)	2 (7)	<0.001
	Fears, worries	34 (62)	0 (0)	<0.001
	Tradeoffs	18 (33)	2 (7)	0.01
	Function, abilities	20 (36)	0 (0)	<0.001

	Quality of life	26 (47)	2 (7)	<0.001
	Family involvement, etc	17 (31)	2 (7)	0.01
End-of-life care planning	End-of-life, unspecified	27 (4)	18 (64)	0.19
	Palliative care, etc	22 (40)	18 (64)	0.04
	Hospice	2 (4)	0 (0)	0.55
	Site of death discussion, etc	7 (13)	0 (0)	0.09

All values reported as number (percent)