

Histone H1 loss drives lymphoma by disrupting 3D chromatin architecture

Nevin Yusufova^{1,2}, Alexey A. Soshnev³, Andreas Kloetgen^{4,5}, Matt Teater¹, Adewola Osunsade⁶, Jeannie M. Camarello⁷, Christopher R. Chin^{8,9}, Ashley Doane^{9,10}, Bryan J. Venters¹¹, Stephanie Portillo-Ledesma¹², Joseph Conway¹³, Jude Philips¹, Olivier Elemento¹⁰, David W. Scott¹⁴, Wendy Beguelin¹, Jonathan D. Licht¹⁵, Neil Kelleher⁷, Louis M Staudt¹⁶, Arthur I. Skoultchi¹⁷, Michael-Christopher Keogh¹¹, Eftychia Apostolou¹⁸, Christopher Mason^{8,9,19}, Marcin Imielinski¹³, Tamar Schlick^{12,19}, Yael David⁶, Aristotelis Tsirigos^{4,20}, C. David Allis³, Ethel Cesarman^{13*}, Ari M. Melnick^{1*}

Supplementary Information Guide

1. Methods
2. Supplementary Figure 1. Uncropped gel blots
3. Supplementary Figure 2. Flow cytometry and sorting gating strategies