

Supplementary Table 3. Comparison of the Sites and Pattern of Vascular Manifestation at Diagnosis between ANCA-Positive and ANCA-Negative BD Patients Based on the 1990 ISG Criteria

Vascular manifestation	Total (n=588)	ANCA-positive BD patients (n=13)	ANCA-negative BD patients (n=575)	p value
Lower extremity vein thrombosis				
Deep vein	6 (1.0)	0 (0)	6 (1.0)	1.000
Superficial vein	5 (0.9)	0 (0)	5 (0.9)	1.000
Vena cava thrombosis				
Inferior vena cava	0 (0)	0 (0)	0 (0)	N/A
Superior vena cava	0 (0)	0 (0)	0 (0)	N/A
Cerebral venous sinus thrombosis	0 (0)	0 (0)	0 (0)	N/A
Pulmonary artery involvement	2 (0.3)	0 (0)	2 (0.3)	1.000
Intracardiac thrombosis	2 (0.3)	1 (7.7)	1 (0.2)	0.044
Aorta involvement				
Aortic aneurysm	3 (0.5)	0 (0)	3 (0.5)	1.000
Carotid artery aneurysm	2 (0.3)	0 (0)	2 (0.3)	1.000
Extra-pulmonary artery involvement				
Lower extremity artery	7 (1.2)	1 (7.7)	6 (1.0)	0.146
Upper extremity artery	2 (0.3)	1 (7.7)	1 (0.2)	0.044
Visceral artery	1 (0.2)	1 (7.7)	0 (0)	0.022
Comments	N=26, E=30	N=3, E=4	N=23, E=26	
Number of patients with vascular manifestation (N) and vascular events (E)				

BD, Behçet's disease; ANCA, antineutrophil cytoplasmic antibody; ISG, International Study Group; E, event.

Values are expressed as number (percentage).