

Supplementary Table S1. Clinical and laboratory data of patients CLL#1 and CLL#5.

Patients CLL#1 and CLL#5 were clinically followed for 2 years, and peripheral blood samples were collected at the indicated time. At the diagnosis, CLL#1 patient counted 98% CD19/CD5 positive B cells, and CLL#5 patient counted 82% CD19/CD5 positive B cells.

Definition of CLL stage was according to Binet classification (Cancer.Net Editorial Board, 10/2017). Cytogenetic analysis was performed using Fluorescent in situ hybridization (FISH) with probes XL p53/LSI ATM and XL DLEU/LAMP/12cen (MetaSystems). WBC, white blood cells.

Patient	Sample (collection time)	WBC	Binet stage	Cytogenetics	Therapy
CLL#1 65-years old male	CLL#1a (month 1)	40,410/mm ³	A	Del13q14	None
	CLL#1b (month 5)	69,070/mm ³	A		None
	CLL#1c (month 8)	92,670/mm ³	C		rituximab-bendamustine (for 6 months)
	CLL#1d (month 24)	6,700/mm ³	A		None
CLL#5 80-years old female	CLL#5a (month 1)	57,210/mm ³	A	Del13q14	None
	CLL#5b (month 12)	119,000/mm ³	A		None
	CLL#5c (month 24)	86,500/mm ³	A		None