

Figure S1. Antifungal activity of compounds 1-13 against FO5C.

*N.T: Not treated (drug-free test).

Compounds that do not share the same letter are statistically different at $P < 0.05$.





Figure S2. Antifungal activity of compounds 1-13 against FSSC.

*N.T: Not treated (drug-free test).

Compounds that did not share the same letter are statistically different at $P < 0.05$.





Figure S3. Antifungal activity of compounds 1-13 against FFSC.

*N.T: Not treated (drug-free test).

Compounds that did not share the same letter are statistically different at $P < 0.05$.






Figure S4. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester **13** (A), TRB (B) and AmB (C) against *F. oxysporum* 89.






Figure S5. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester 13 (A), TRB (B) and AmB (C) against *F. oxysporum* 91.






Figure S6. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester 13 (A), TRB (B) and AmB (C) against *F. keratoplasicum* 93.






Figure S7. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester **13** (A), TRB (B) and AmB (C) against *F. solani* 96.






Figure S8. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester 13 (A), TRB (B) and AmB (C) against *F. verticillioides* 87.




Figure S9. MIC and LD₅₀ ranges expressed as absorbance (milliOD) at 595 nm at 48 h of Ester 13 (A), TRB (B) and AmB (C) against *F. verticillioides* 115.