

**S2 Table. Adjusted analysis of Choosing Wisely recommendation 3.**

(3) Discuss whether treatment limitations are needed when talking to patients about treatment options

<b>Variable</b>	<b>Beta intervention*</b>	<b>Bivariable OR (95% CI)</b>	<b>P value</b>	<b>Multivariable OR (95% CI)</b>	<b>P value</b>
Intervention*	0.340	1.405 (0.663–2.978)	0.374	1.431 (0.659–3.107)	0.365
Age	0.354	0.995 (0.975–1.015)	0.602		
Male Sex	0.333	0.390 (0.182–0.835)	0.015		
Hospital	0.282	3.967 (1.799–8.751)	0.001	4.610 (2.060–10.316)	<0.001
Specialism	0.336	1.015 (0.789–1.306)	0.906		
Charlson comorbidity index	0.347	1.092 (0.509–2.343)	0.821		
High MEWS score	0.326	0.834 (0.240–2.891)	0.774		
Hospitalized 3 mo before ED visit	0.387	0.390 (0.155–0.980)	0.045	0.308 (0.120–0.793)	0.015
Outpatient care for internal medicine 1 year before presentation at the ED	0.354	0.424 (0.196–0.919)	0.030		
Nursing home resident	0.341	0.841 (0.107–6.638)	0.870		