

OMTO, Volume 20

Supplemental Information

**Genetic variants in m⁶A modification
core genes are associated
with glioma risk in Chinese children**

Jing He, Li Yuan, Huiran Lin, Ao Lin, Huitong Chen, Ailing Luo, Zhenjian Zhuo, and Xiaoping Liu

Table S1. Frequency distribution of selected variables in glioma patients and cancer-free controls

Variables	Cases (N=171)		Controls (N=228)		<i>P</i> ^a
	No.	%	No.	%	
Age range, month	4.00-168.00		4.00-168.00		0.623
Mean ± SD	63.40 ± 47.72		52.41 ± 32.65		
<60	85	49.71	119	52.19	0.190
≥60	86	50.29	109	47.81	
Gender					0.190
Female	81	47.37	93	40.79	
Male	90	52.63	135	59.21	
Subtypes					
Astrocytic tumors	125	73.10	/	/	
Ependymoma	24	14.62	/	/	
Neuronal and mixed neuronal-gliial tumours	14	8.19	/	/	
Embryonal tumors	7	4.09	/	/	
Tumor grades					
I	103	60.23	/	/	
II	28	16.37	/	/	
III	15	8.77	/	/	
IV	25	14.62	/	/	

SD, standard deviation.

^a Two-sided χ^2 test for distributions between glioma patients and cancer-free controls.