
a
M-cutotype
C-cutotype
C-cutotype enriched species
M-cutotype enriched species
Cutibacterium sp.
Lawsonella clevelandensis
Staphylococcus sp.
Propionibacterium phage
Staphylococcus phage
Listeria monocytogenes
Jonesia denitrificans
Corynebacterium sp.
Auricoccus indicus
Acinetobacter gyllenbergii
Moraxella sp.
Psychrobacter sp.
Elizabethkingia sp.
Acinetobacter sp.
Chryseobacterium sp.
Paracoccus sp.
Brevundimonas sp.
Deinococcus sp.
Microbacterium sp.
Novosphingobium sp.
Rhodobacter sp.
high
low
b
Malassezia globosa
Malassezia slooffiae
Corynebacterium sp.
Cutibacterium acnes
Cutibacterium granulosum
Moraxella osloensis
c
Staphylococcus capitis
Propionibacterium sp.
Staphylococcus sciuri
Cutibacterium acnes
Moraxella osloensis
Staphylococcus epidermidis
d
Staphylococcus capitis
Bacillus subtilis
Staphylococcus sciuri
Cutibacterium acnes
Streptococcus suis
Streptococcus salivarius
Americans
e
R² = 0.3767
P value = 0.001
PCoA1 47 %
PCoA2 10.29 %
Jensen–Shannon
R² = 0.4831
P value = 0.001
PCoA1: 58.94 %
PCoA2: 11.81 %
Bray-Curtis
f
R² = 0.3014
P value = 0.001
PCoA1 39.57 %
PCoA2 11.79 %
Jensen–Shannon
R² = 0.3942
P value = 0.001
PCoA1: 51.64 %
PCoA2: 13.58 %
Bray-Curtis