

Supplementary Table 1. Patients' characteristics by IV tPA status

Characteristic	IV tPA (n=28)	No IV tPA (n=66)	P
Baseline NIHSS	2.04±1.71	2.15±1.61	0.755
Onset-to-door time (hr)	1.33 (0.89–2.20)	5.36 (2.38–13.38)	<0.001
Site of occlusion			0.134
Intracranial ICA	0 (0)	7 (10.6)	
Tandem	1 (3.6)	5 (7.6)	
MCA M1	12 (42.9)	32 (48.5)	
MCA M2 trunk	15 (53.6)	22 (33.3)	
Baseline glucose (mg/dL)	144.2±53.5	143.6±50.9	0.963
Initial DWI volume (mL)	0.52 (0.00–5.97)	1.53 (0.17–5.52)	0.380
Tmax >6 sec (mL)	40.62 (15.86–57.96)	25.47 (5.23–52.30)	0.383
Tmax >6 sec – DWI (mL)	37.42 (12.36–54.64)	21.79 (3.18–41.37)	0.298
Tmax 4–6 sec (mL)	35.09 (26.34–83.93)	51.16 (28.42–81.39)	0.185

Values are presented as mean±standard deviation, median (interquartile range), or number (%).

IV tPA, intravenous tissue-type plasminogen activator; NIHSS, National Institutes of Health Stroke Scale; ICA, internal carotid artery; MCA, middle cerebral artery; DWI, diffusion-weighted imaging.