

Supplementary Table 3. Imaging variables of enrolled patients and patients who underwent EVT before END

Variable	Enrolled patients (n=94)	Excluded patients: EVT before END*	P
Initial DWI volume (mL)	1.34 (0.00–5.61)	3.45 (1.46–7.80)	0.065
Tmax >2 sec (mL)	217.62 (156.56–274.25)	273.00 (215.14–357.00)	0.044
Tmax >4 sec (mL)	84.24 (47.78–120.57)	128.65 (77.16–208.59)	0.013
Tmax >6 sec (mL)	28.97 (10.15–56.67)	62.33 (28.81–131.16)	0.018
Tmax >8 sec (mL)	12.53 (0.00–25.10)	32.46 (3.47–77.81)	0.080
Tmax >10 sec (mL)	3.85 (0.00–12.70)	16.41 (0.00–45.91)	0.082
Tmax >4 sec – DWI (mL)	75.21 (43.27–118.17)	128.65 (72.89–206.64)	0.017
Tmax >6 sec – DWI (mL)	24.96 (6.45–46.25)	51.98 (25.61–129.21)	0.024
Tmax 2–6 sec (mL)	185.54 (131.62–229.11)	198.23 (154.99–233.46)	0.709
Tmax 4–6 sec (mL)	47.19 (28.10–76.63)	60.11 (45.70–76.88)	0.353

Values are presented as median (interquartile range).

EVT, endovascular therapy; END, early neurological deterioration; DWI, diffusion-weighted imaging.

*Of the 14 excluded patients who received EVT before END, those who had missing data on DWI (n=1) and perfusion variables (n=5) were excluded from the corresponding analysis.