

Supplementary Table S1. Baseline and follow-up results of non-surgical periodontal therapy for study population of part 2 ($n = 91$).

| | Baseline | 3 mo | 6 mo | 12 mo |
|----------------------|-----------------|-------------|-------------|--------------|
| Mean PPD (mm) | 3.9 ± 0.6 | 2.8 ± 0.4 | 2.8 ± 0.4 | 2.8 ± 0.5 |
| Mean CAL (mm) | 4.2 ± 0.9 | 3.7 ± 1.0 | 3.6 ± 1.0 | 3.6 ± 1.0 |
| Mean BOP (%) | 66.3 ± 14.8 | 21.2 ± 12.7 | 20.7 ± 13.5 | 21.8 ± 13.8 |
| Mean plaque (%) | 62.8 ± 24.2 | 18.2 ± 16.4 | 21.7 ± 20.2 | 20.1 ± 15.6 |
| Sites PPD ≥ 6 mm (%) | 34.0 ± 17.7 | 6.8 ± 7.5 | 6.2 ± 7.9 | 6.5 ± 8.2 |

Clinical parameters of patients **of part 2** before (Baseline) and 3, 6, and 12 mo after non-surgical treatment. This table is adjusted from the previously published study by Bizzarro et al. (2016). Values represent means ± SD.