

**S1 Table. Histopathologic findings in AGMs and RMs following inhalation of *B. pseudomallei* HBPUB10134a.**

	Prevalance		Severity <sup>a</sup> : average (range)	
	AGM	RM	AGM	RM
<b>Pneumonia</b>	5/5	3/5	3.8 (3-5)	3.3 (2-4)
<b>Pleuritis</b>	4/5	2/5	3.7 (3-5)	3 (2-4)
<b>Pulmonary pyogranulomas</b>	0/5	1/5	N/A	4 (4)
<b>Hepatitis</b>	5/5	3/5	2.4 (2-3)	2 (2-3)
<b>Splenitis</b>	5/5	4/5	4.4 (3-4)	2.5 (2-3)
<b>Lymphadenitis</b>				
<b>Tracheobronchial</b>	3/4 <sup>b</sup>	2/5	2.6 (2-3)	2 (2)
<b>Mediastinal</b>	4/5	1/5	3.2 (3-4)	2 (2)
<b>Nephritis</b>	0/5	2/5	N/A	1.5 (1-2)
<b>Myelitis (bone marrow)</b>	1/5	0/5	1 (1)	N/A
<b>Bone marrow pyogranulomas</b>	4/5	0/5	2.5 (2-4)	N/A

<sup>a</sup>Severity scores: 1: minimal, 2: mild, 3: moderate, 4: marked, 5: severe.

<sup>b</sup>One tracheobronchial lymph node was not examined.