Figure 1b



Figure 1c



Figure 1d



Figure 1e



Figure 1g



Figure 2a



Same gel (PS1-NTF: 800 exposure, β-Actin: 680 exposure)

Figure 2c



Same gel (Top: 680 exposure, Bottom: 800 exposure)

Figure 3a



Same gel
(PS1-NTF: 800 exposure,
β-Actin: 680 exposure)

Figure 3c



Figure 3e

Same sample prep, separate gel (β-Actin)

Figure 3d



Same gel (Top: 680 exposure, Middle: 800 exposure)

Figure 3f



Same gel (Top: 800 exposure, Bottom: 680 exposure)



Separate gel

Figure 4c



Separate gel (β-Actin, * : no sample)

Figure 4e



Same gel (β-Tubulin III)

Figure 4h



Figure 4i



Same gel (Left: 680 exposure, Right: 800 exposure)



Figure 5b



Figure 5c



Figure 5e



Extended Data Figure 1b



Extended Data Figure 1d

Extended Data Figure 1f



Extended Data Figure 1g



Same gel (Left: 680 exposure, Right: 800 exposure)

Extended Data Figure 1h



Same gel (Tubulin)

Extended Data Figure 2g



Same gel (β-Actin: 680 exposure, NICD (Val1744): 800 exposure)

Extended Data Figure 2h



Extended Data Figure 3b



Extended Data Figure 4k



Same gel (β-Tubulin III)

Extended Data Figure 4m



Extended Data Figure 5b