

Mandibular Advancement Device Treatment Efficacy is Associated with Polysomnographic Endotypes

Online Supplement

Sara Op de Beeck MSc^{*1, 3, 6}, Marijke Dieltjens PhD^{1, 5, 6}, Ali Azarbarzin PhD², Marc Willemen MSc³, Johan Verbraecken PhD^{1, 3, 4}, Marc J. Braem PhD^{1, 5}, Andrew Wellman PhD², Scott A. Sands PhD^{2°}, Olivier M. Vanderveken PhD^{1, 3, 6°}

°Equal contributions

¹ Faculty of Medicine and Health Sciences, University of Antwerp, Wilrijk, Belgium

² Division of Sleep and Circadian Disorders, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, United States,

³ Multidisciplinary Sleep Disorders Centre, Antwerp University Hospital, Edegem, Belgium,

⁴ Department of Pulmonology, Antwerp University Hospital, Edegem, Antwerp, Belgium,

⁵ Special Dentistry Care, Antwerp University Hospital, Edegem, Belgium,

⁶ ENT, Head and Neck Surgery, Antwerp University Hospital, Edegem, Belgium

*Corresponding author:

Sara Op de Beeck
Translational Neurosciences
Faculty of Medicine and Health Sciences
University of Antwerp
Universiteitsplein 1 – D.T.494
2610 Wilrijk
BELGIUM
Email: sara.opdebeeck@uantwerpen.be

Supplemental Figures




Figure E1: Study protocol.

PSG: polysomnography, MAD: mandibular advancement device, AHI: apnea-hypopnea index