

Table ESM 3: *Future years of life lost (1000s) among currently prevalent diabetes patients in Denmark at 1 January each year, and average years of life lost per person among these. Note that only every 3 January is shown.*

	Date	Total YLL (1000s)			Average YLL	
		T1D	T2D	DM	T1D	T2D
Men	1996	85.1	149.4	234.4	6.9	4.9
	1999	110.9	188.2	299.2	8.4	4.6
	2002	129.8	222.8	352.6	9.5	4.3
	2005	138.8	266.8	405.6	9.9	3.9
	2008	148.1	276.8	424.9	10.3	3.4
	2011	141.4	310.5	451.8	9.6	3.1
	2014	121.2	377.9	499.1	8.0	3.0
	2017	99.9	428.5	528.4	6.4	3.1
Women	1996	74.7	150.6	225.3	7.8	4.8
	1999	86.6	173.5	260.2	8.6	4.4
	2002	94.8	192.6	287.3	9.1	4.1
	2005	102.2	223.3	325.5	9.7	3.7
	2008	108.5	225.4	334.0	10.0	3.2
	2011	103.2	248.4	351.7	9.3	3.0
	2014	92.7	303.7	396.3	8.1	2.9
	2017	81.9	337.3	419.2	6.9	3.0
M+W	1996	159.8	300.0	459.7	7.3	4.9
	1999	197.6	361.8	559.3	8.5	4.5
	2002	224.6	415.4	639.9	9.3	4.2
	2005	241.0	490.1	731.1	9.8	3.8
	2008	256.7	502.2	758.8	10.2	3.3
	2011	244.6	558.9	803.5	9.5	3.0
	2014	213.9	681.5	895.4	8.0	3.0
	2017	181.8	765.8	947.6	6.6	3.0