

Supplemental material**Table 1.1: Details of the studies included in targeted literature review for the three UBT devices**

Authors	Study design	PPH success rate	Atonic PPH success rate	Reference
Condom-UBT				
Darwish et al.	RCT	28/33 (84.8)	28/33 (84.8)	[1]
Tindell et al.	Systematic Review	186/193 (96.4)	NR	[2]
Santhanam et al.	Prospective	59/61 (96.7)	59/61 (96.7)	[3]
Rathore et al.	Prospective	17/18 (94.4)	NR	[4]
Aderoba et al.	Prospective	203/229 (88.6)	193/214 (90.2)	[5]
Mishra et al.	Prospective	59/60 (98.3)	NR	[6]
Kandeel et al.	Prospective	48/50 (96.0)	28/28 (100)	[7]
Anger et al.	RCT	56/64 (87.5)	NR	[8]
Dumont et al.	RCT	48/57 (84.2)	NR	[9]
Lohano et al.	Prospective	126/139 (90.6)	126/139 (90.6)	[10]
Hasabe et al.	Prospective	34/36 (94.4)	NR	[11]
Yadav et al.	Prospective	117/122 (95.9)	117/122 (95.9)	[12]
Bakri-UBT				
Darwish et al.	RCT	30/33 (90.9)	30/33 (90.9)	[1]
Revert et al.	Prospective	188/226 (83.2)	155/183 (84.7)	[13]
Brown et al.	Prospective	55/58 (94.8)	52/55 (94.5)	[14]
Vintejoux et al.	Retrospective	25/36 (69.4)	25/36 (69.4)	[15]
Guo et al.	Retrospective	288/305 (94.4)	131/142 (92.3)	[16]
Mathur et al.	Retrospective	40/49 (81.6)	14/17 (82.4)	[17]
Wang et al.	Prospective	373/407 (91.6)	373/407 (91.6)	[18]
Alkis et al.	Retrospective	43/47 (91.5)	NR	[19]
Kaya et al.	Prospective	34/45 (75.6)	27/34 (79.4)	[20]
Laas et al.	Before and after	37/43 (86)	37/43 (86)	[21]
Olsen et al.	Retrospective	25/37 (67.6)	17/24 (70.8)	[22]
Kong et al.	Retrospective	59/81 (72.8)	37/59 (62.7)	[23]

Cetin et al.	Retrospective	29/39 (74.4)	29/39 (74.4)	[24]
Gauchotte et al.	Before and after	35/38 (92.1)	NR	[25]
Grange et al.	Retrospective	80/108 (74.1)	26/39 (66.7)	[26]
Kadioglu et al.	Retrospective	42/50 (84)	NR	[27]
Martin et al.	Retrospective	32/49 (65.3)	28/42 (66.7)	[28]
Ogoyama et al.	Retrospective	66/71 (93)	31/32 (96.9)	[29]
Son et al.	Retrospective	239/306 (78.1)	190/241 (78.8)	[30]
ESM-UBT				
Ramanathan et al	Prospective/ Retrospective case series	189/201 (94) *	NR	[31]
Burke et al.	Prospective case series	190/201 (94.5) *	NR	[32]
Burke et al.	Prospective case series	298/306 (97.4) *	298/306 (97.4)	[33]

* - Reported survival rates

NR – Not reported

REFERENCES

- 1 Darwish AM, Abdallah MM, Shaaban OM, et al. Bakri balloon versus condom-loaded Foley's catheter for treatment of atonic postpartum hemorrhage secondary to vaginal delivery: a randomized controlled trial. *J Matern Neonatal Med* 2018;**31**:747–53.
doi:10.1080/14767058.2017.1297407
- 2 Tindell K, Garfinkel R, Abu-Haydar E, et al. Uterine balloon tamponade for the treatment of postpartum haemorrhage in resource-poor settings: A systematic review. *BJOG An Int J Obstet Gynaecol*. 2013;**120**:5–14. doi:10.1111/j.1471-0528.2012.03454.x
- 3 Santhanam R, Viswanathan RM, V. P. Condom tamponade in the management of atonic postpartum hemorrhage. *Int J Reprod Contraception, Obstet Gynecol* 2018;**7**:2276.
doi:10.18203/2320-1770.ijrcog20182335
- 4 Rathore AM, Gupta S, Manaktala U, et al. Uterine tamponade using condom catheter balloon in the management of non-traumatic postpartum hemorrhage. *J Obstet Gynaecol Res* 2012;**38**:1162–7. doi:10.1111/j.1447-0756.2011.01843.x
- 5 Aderoba A, Olagbuji B, Akintan A, et al. Condom-catheter tamponade for the treatment of postpartum haemorrhage and factors associated with success: a prospective observational study. *BJOG An Int J Obstet Gynaecol* 2017;**124**:1764–71. doi:10.1111/1471-0528.14361
- 6 Mishra N, Gulabani K, Agrawal S, et al. Efficacy and Feasibility of Chhattisgarh Balloon and Conventional Condom Balloon Tamponade: A 2-Year Prospective Study. *J Obstet Gynecol India* 2019;**69**:133–41. doi:10.1007/s13224-018-1185-6
- 7 Kandeel M, Sanad Z, Ellakwa H, et al. Management of postpartum hemorrhage with intrauterine balloon tamponade using a condom catheter in an Egyptian setting. *Int J Gynecol Obstet* 2016;**135**:272–5. doi:10.1016/j.ijgo.2016.06.018
- 8 Anger HA, Dabash R, Durocher J, et al. The effectiveness and safety of introducing condom-catheter uterine balloon tamponade for postpartum haemorrhage at secondary level hospitals in Uganda, Egypt and Senegal: a stepped wedge, cluster-randomised trial. *BJOG An Int J Obstet Gynaecol* 2019;**126**:1612–21. doi:10.1111/1471-0528.15903
- 9 Dumont A, Bodin C, Hounkpatin B, et al. Uterine balloon tamponade as an adjunct to misoprostol for the treatment of uncontrolled postpartum haemorrhage: A randomised controlled trial in Benin and Mali. *BMJ Open* 2017;**7**:1–9. doi:10.1136/bmjopen-2017-016590
- 10 Lohano R, Haq G, Kazi S, et al. Intrauterine balloon tamponade for the control of postpartum haemorrhage. *J Pak Med Assoc* 2016;**66**:22–6.
- 11 Hasabe R, Gupta K, Rathode P. Use of Condom Tamponade to Manage Massive Obstetric Hemorrhage at a Tertiary Center in Rajasthan. *J Obstet Gynecol India* 2016;**66**:88–93.
doi:10.1007/s13224-015-0790-x
- 12 Yadav S, Malhotra A. A prospective randomized comparative study of Misoprostol and balloon tamponade using condom catheter to prevent postpartum hemorrhage at M. Y. H., Indore, India in vaginal delivered patients. *Int J Reprod Contraception, Obstet Gynecol* 2019;**8**:591. doi:10.18203/2320-1770.ijrcog20190290
- 13 Revert M, Cottenet J, Raynal P, et al. Intrauterine balloon tamponade for management of severe postpartum haemorrhage in a perinatal network: a prospective cohort study. *BJOG An Int J Obstet Gynaecol* 2017;**124**:1255–62. doi:10.1111/1471-0528.14382
- 14 Brown H, Okeyo S, Mabeya H, et al. The Bakri tamponade balloon as an adjunct treatment for refractory postpartum hemorrhage. *Int J Gynecol Obstet* 2016;**135**:276–80.
doi:10.1016/j.ijgo.2016.06.021
- 15 Vintejoux E, Ulrich D, Mousty E, et al. Success factors for Bakri™ balloon usage secondary to uterine atony: A retrospective, multicentre study. *Aust New Zeal J Obstet Gynaecol* 2015;**55**:572–7. doi:10.1111/ajo.12376
- 16 Guo Y, Hua R, Bian S, et al. Intrauterine Bakri Balloon and Vaginal Tamponade Combined with Abdominal Compression for the Management of Postpartum Hemorrhage. *J Obstet Gynaecol Canada* 2018;**40**:561–5. doi:10.1016/j.jogc.2017.08.035
- 17 Mathur M, Ng QJ, Tagore S. Use of Bakri balloon tamponade (BBT) for conservative management of postpartum haemorrhage: a tertiary referral centre case series. *J Obstet*

- 18 *Gynaecol (Lahore)* 2018;**38**:66–70. doi:10.1080/01443615.2017.1328671
18 Wang D, Xu S, Qiu X, et al. Early usage of Bakri postpartum balloon in the management of postpartum hemorrhage: A large prospective, observational multicenter clinical study in South China. *J Perinat Med* 2018;**46**:649–56. doi:10.1515/jpm-2017-0249
- 19 Alkiş I, Karaman E, Han A, et al. The fertility sparing management of postpartum hemorrhage: A series of 47 cases of Bakri balloon tamponade. *Taiwan J Obstet Gynecol* 2015;**54**:232–5. doi:10.1016/j.tjog.2014.03.009
- 20 Kaya B, Tuten A, Daglar K, et al. Balloon tamponade for the management of postpartum uterine hemorrhage. *J Perinat Med* 2014;**42**:745–53. doi:10.1515/jpm-2013-0336
- 21 Laas E, Bui C, Popowski T, et al. Trends in the rate of invasive procedures after the addition of the intrauterine tamponade test to a protocol for management of severe postpartum hemorrhage. *Am J Obstet Gynecol* 2012;**207**:281.e1–281.e7. doi:10.1016/j.ajog.2012.08.028
- 22 Olsen R, Reisner DP, Benedetti TJ, et al. Bakri balloon effectiveness for postpartum hemorrhage: A ‘real world experience’. *J Matern Neonatal Med* 2013;**26**:1720–3. doi:10.3109/14767058.2013.796354
- 23 Kong CW, To WW. Prognostic factors for the use of intrauterine balloon tamponade in the management of severe postpartum hemorrhage. *Int J Gynecol Obstet* 2018;**142**:48–53. doi:10.1002/ijgo.12498
- 24 Çetin BA, Aydogan Mathyk B, Atis Aydin A, et al. Comparing success rates of the Hayman compression suture and the Bakri balloon tamponade. *J Matern Neonatal Med* 2019;**32**:3034–8. doi:10.1080/14767058.2018.1455184
- 25 Gauchotte E, De La Torre M, Perdriolle-Galet E, et al. Impact of uterine balloon tamponade on the use of invasive procedures in severe postpartum hemorrhage. *Acta Obstet Gynecol Scand* 2017;**96**:877–82. doi:10.1111/aogs.13130
- 26 Grange J, Chatellier M, Chevé MT, et al. Predictors of failed intrauterine balloon tamponade for persistent postpartum hemorrhage after vaginal delivery. *PLoS One* 2018;**13**:1–11. doi:10.1371/journal.pone.0206663
- 27 Kadioglu BG, Tanrıverdi EC, Aksoy AN. Balloon Tamponade in the Management of Postpartum Hemorrhage: Three Years of Experience in a Single Center. *Open J Obstet Gynecol* 2016;**06**:698–704. doi:10.4236/ojog.2016.612087
- 28 Martin E, Legendre G, Bouet PE, et al. Maternal outcomes after uterine balloon tamponade for postpartum hemorrhage. *Acta Obstet Gynecol Scand* 2015;**94**:399–404. doi:10.1111/aogs.12591
- 29 Ogoyama M, Takahashi H, Usui R, et al. Hemostatic effect of intrauterine balloon for postpartum hemorrhage with special reference to concomitant use of “holding the cervix” procedure (Matsubara). *Eur J Obstet Gynecol Reprod Biol* 2017;**210**:281–5. doi:10.1016/j.ejogrb.2017.01.012
- 30 Son M, Einerson BD, Schneider P, et al. Is There an Association between Indication for Intrauterine Balloon Tamponade and Balloon Failure? *Am J Perinatol* 2017;**34**:164–8. doi:10.1055/s-0036-1585084
- 31 Ramanathan A, Eckardt MJ, Nelson BD, et al. Safety of a condom uterine balloon tamponade (ESM-UBT) device for uncontrolled primary postpartum hemorrhage among facilities in Kenya and Sierra Leone. *BMC Pregnancy Childbirth* 2018;**18**:1–7. doi:10.1186/s12884-018-1808-z
- 32 Burke TF, Ahn R, Nelson BD, et al. A postpartum haemorrhage package with condom uterine balloon tamponade: a prospective multi-centre case series in Kenya, Sierra Leone, Senegal, and Nepal. *BJOG An Int J Obstet Gynaecol* 2016;**123**:1532–40. doi:10.1111/1471-0528.13550
- 33 Burke TF, Danso-Bamfo S, Guha M, et al. Shock progression and survival after use of a condom uterine balloon tamponade package in women with uncontrolled postpartum hemorrhage. *Int J Gynecol Obstet* 2017;**139**:34–8. doi:10.1002/ijgo.12251
- 34 Joseph CM, Bhatia G, Abraham V, et al. Obstetric admissions to tertiary level intensive care unit – Prevalence, clinical characteristics and outcomes. *Indian J Anaesth* 2018;**62**:940–4. doi:10.4103/ija.IJA_537_18
- 35 Rastogi A, Zahid. Postpartum haemorrhage | National Health Portal Of India. Natl. Heal. Portal India. 2017. <https://www.nhp.gov.in/disease/gynaecology-and-obstetrics/postpartum-haemorrhage/>

- haemorrhage (accessed 20 Jul 2020).
- 36 Carroli G, Cuesta C, Abalos E, et al. Epidemiology of postpartum haemorrhage: a systematic review. *Best Pract. Res. Clin. Obstet. Gynaecol.* 2008;22:999–1012.
doi:10.1016/j.bpobgyn.2008.08.004
- 37 Dutta D. *Textbook of Obstetrics*. 2015.
- 38 Tasneem F, Sirsam S, Shanbhag V. Clinical study of post partum haemorrhage from a teaching hospital in Maharashtra, India. *Int J Reprod Contraception, Obstet Gynecol* 2017;6:2366.
doi:10.18203/2320-1770.ijrcog20172314
- 39 International Institute for Population Sciences. National Family Health Survey (NFHS-4) 2015-16 India. *Int Inst Popul Sci ICF* 2017;;1–192. doi:kwm120 [pii]10.1093/aje/kwm120