



Prevalence, Predictors, and Mid-Term Outcomes of Non-Home Discharge After Transcatheter Aortic Valve Implantation

Yuuki Muto, MD; Ryosuke Higuchi, MD; Akiomi Yoshihisa, MD, PhD;
Kenichi Hagiya, MD; Mike Saji, MD; Itaru Takamisawa, MD;
Tetsuya Tobaru, MD, PhD; Nobuo Iguchi, MD, PhD; Shuichiro Takanashi, MD, PhD;
Morimasa Takayama, MD, PhD; Yasuchika Takeishi, MD, PhD; Mitsuaki Isobe, MD, PhD

Supplementary Table. The Definitions of Patient Comorbidities

Comorbidity	Definition
Hypertension	The recent use of antihypertensive drug
Diabetes mellitus	The recent use of insulin or antidiabetic drug
Peripheral artery disease	ABI less than 0.9 or previous revascularization of lower limb
Previous stroke	Previous symptomatic cerebrovascular event including TIA, cerebral infarction and cerebral hemorrhage
AF/AFL	The documentation on ECG or medical history on patient record
COPD	FEV ₁ % less than 70% or the recent use of drug for COPD
OMI	At least one or more finding compatible with myocardial infarction on ECG, echocardiography, radioisotope imaging or MRI
Previous CABG	Previous isolated or concurrent CABG

ABI, ankle-brachial index; TIA, transient ischemic attack; AF, atrial fibrillation; AFL, atrial flutter; ECG, electrocardiogram; COPD, chronic obstructive pulmonary disease; FEV, forced expiratory volume; OMI, old myocardial infarction; MRI, magnetic resonance imaging; CABG, coronary artery bypass grafting.