A Multicenter, Prospective, Observational Study to Determine Association of Mesangial C1q Deposition with Renal Outcomes in IgA Nephropathy

Li Tan, MD ^{1,5,6}, Yi Tang, MD ¹, Gaiqin Pei, MD ^{1,6}, Zhengxia Zhong, MD ^{2,6}, Jiaxing Tan, MD ^{1,6}, Ling Zhou, MD ^{3,6}, Dongmei Wen, MD ^{4,6}, David Sheikh-Hamad, MD ⁵, Wei Qin, MD ¹

¹ Division of Nephrology, Department of Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, China.

² Division of Nephrology, Department of Medicine, Affiliated Hospital of Zunyi Medical University, Medical University, Zunyi, Guizhou, China.


³ Division of Nephrology, Zigong Third People's Hospital, Zigong, Sichuan, China.

⁴ Division of Nephrology, People's Hospital of Jianyang, Chengdu, Sichuan, China.

⁵ Section of Nephrology, Department of Medicine, Baylor College of Medicine, Houston, TX, USA.

⁶ West China School of Medicine, Sichuan University, Chengdu, Sichuan, China.

Correspondence to: Wei Qin, Division of Nephrology, Department of Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, China. Tel. 86-28-85422338, Fax +86-028-8542-3341. Email qinweihx@scu.edu.cn.



Supplementary Fig. S1. Treatment response and renal outcome of IgAN patients.

Abbreviations: CR, complete remission; PR, partial remission; NR, no response; ESRD, end stage renal disease.