

Supplementary Table S1. Results from crude and weighted models for the association between past year PM_{2.5} (10 µg/m³ increase) with obesity in children, adolescents and adults* from the Greater Mexico City Area using data from the National Nutrition and Health Surveys, ENSANUT-2006 and 2012.

	Children		Adolescents		Adults	
	OR	95% CI	OR	95% CI	OR	95% CI
ENSANUT 2006						
<i>Weighted model</i>	<i>n</i> = 2,529,289		<i>n</i> = 3,548,352		<i>n</i> = 12,541,729	
	1.24	(0.49, 3.13)	3.53	(1.45, 8.58)	1.01	(0.59, 1.73)
<i>Crude model</i>	<i>n</i> = 618		<i>n</i> = 801		<i>n</i> = 1559	
	0.95	(0.39, 2.28)	2.38	(1.05, 5.38)	1.09	(0.66, 1.81)
ENSANUT 2012						
<i>Weighted model</i>	<i>n</i> = 2,769,354		<i>n</i> = 3,245,296		<i>n</i> = 11,415,512	
	1.77	(0.82, 3.86)	3.79	(1.40, 10.24)	2.73	(0.97, 7.71)
<i>Crude model</i>	<i>n</i> = 753		<i>n</i> = 758		<i>n</i> = 1559	
	1.96	(0.56, 6.77)	2.44	(0.76, 7.84)	2.16	(1.04, 4.48)

Statistically significant results (p <0.05) are highlighted in bold letters.