

On-line Table 1: Outlier CL and WML analysis for each subgroup of patients with RRMS^a

	RRMS-NI				RRMS-I			
	Standard Lesions		Outliers		Standard Lesions		Outliers	
	WML	CL	WML	CL	WML	CL	WML	CL
CBF (mL/100 g per min)	14.2 (9.6–19.5)	21.0 (14.3–30.3)	46.2 (39.0–57.2)	78.8	12.1 (8.0–17.7)	22.5 (15.3–31.0)	44.5 (38.0–60.6)	64.2 (63.9–64.5)
CBV (mL/100 g)	1.1 (0.8–1.5)	1.3 (1.1–1.9)	3.1 (2.8–3.6)	4.6	1.0 (0.7–1.4)	1.6 (1.2–2.3)	3.3 (2.9–3.8)	5.2
MTT (min)	4.8 (4.1–5.9)	4.3 (3.5–4.8)	0.9 (–23.6 to 8.7)	8.2 (7.8–8.5)	5.1 (4.3–6.0)	4.6 (3.9–5.4)	–1.3 (–269.5 to 1.0)	NA

Note:—NA indicates not applicable.

^a All values are medians, and IQRs are in parentheses. Cortical lesions and white matter lesions have been divided into standard lesions (lesions that are not outliers) and outliers.

On-line Table 2: Dispersion of WMLs and CLs for each subgroup of patients with RRMS^a

	RRMS-NI		RRMS-I	
	CL	WML	CL	WML
CBF	3.6	4.8	3.9	4.8
CBV	3.4	5.1	4.1	5.6
MTT	3.5	5.2	3.9	5.5

^a CBF is in milliliters/100 g per minute; CBV is in milliliters/100 g.