

On-line Table: Individual radiologic data of the 12 infants, 8 without and 4 with arthrogryposis, included in the study^a

No.	Sex	Age at MRI (day)	Smallest AP Diameters (mm)			Largest AP Diameters (mm)			Vertebral Canal AP Diameter (mm)			Conus Medullaris Nerve Roots	Grade of Nerve Root Reduction	Congenital Hip Dysplasia
			Cervical	High Thoracic	Conus Medullaris Base	Cervical	Lumbar	Cervical	Thoracic	Lumbar	Thoracic			
Congenital Zika syndrome without arthrogryposis														
1	F	129	5.4	4.3	6.5	5.5	7.1	9.4	8.8	11.6	A < P	Mild	Yes	
2	M	133	4.6	3.7	5.6	5.3	7.5	11.9	8.7	13.1	A < P	Moderate	Yes	
3	F	157	5.3	4.2	6.5	5.6	7.6	10.1	12.2	13.6	A < P	Mild	Yes	
4	M	139	6.3	3.6	5.9	6.4	7.4	10.3	10.8	13.1	Normal	—	No	
5	M	173	5.7	4.5	6.9	5.7	7.4	9.9	10.2	11.3	A < P	Mild	Yes	
6	F	138	5.1	4.6	6.3	5.7	7.1	9.1	9.6	11.5	Normal	—	Yes	
7	M	113	3.9	3.4	4.6	4.1	5.4	9.5	11.0	11.1	A < P	Moderate	No	
8	M	123	5.9	3.7	6.2	7.0	7.2	9.7	10.2	11.0	A < P	Mild	Yes	
Congenital Zika syndrome with arthrogryposis														
9	F	135	5.0	3.5	5.0	5.1	6.0	10.8	8.4	11.4	A < P	Severe	Yes	
10	M	148	4.4	2.6	3.3	4.6	3.7	9.4	9.9	12.5	ANV	Severe	Yes	
11	M	137	4.5	4.4	4.0	4.5	4.7	11.2	11.5	11.5	A < P	Severe	Yes	
12	F	105	4.5	3.6	4.6	4.8	5.3	10.7	8.9	11.1	ANV	Severe	Yes	

Note:—A indicates anterior root; P, posterior root; AP, anteroposterior; NV, not visualized.

^a Quantitative data are diameter measures (mm).