


ON-LINE FIG 1. Flow diagram of patients in the study.



ON-LINE FIG 2. Histogram of propensity scores.

On-line Table: Sensitivity analyses results for infarction

	OR (95% CI)	P Value
A. Assumes patients with no follow-up imaging did not have an infarction		
Frontal lobe IPH after treatment	3.1 (1.0–10.7)	.05
Non-vasospasm ischemic frontal lobe infarction	9.0 (2.0–66.2)	.004
Artery of Heubner infarction after treatment	22.2 (3.7–431.2)	<.001
Embolic infarction after treatment	0.3 (0.1–1.1)	.08
Vasospasm-related ischemic infarction	0.9 (0.3–3.2)	.90
Any ischemic infarction after treatment	4.3 (1.6–12.5)	.004
B. Assumes patients with no follow-up imaging had an infarction		
Frontal lobe IPH after treatment	2.2 (0.8–6.2)	.13
Non-vasospasm ischemic frontal lobe infarction	3.9 (1.2–16.1)	.03
Artery of Heubner infarction after treatment	6.5 (1.8–31.6)	.003
Embolic infarction after treatment	0.4 (0.1–1.1)	.07
Vasospasm-related ischemic infarction	0.8 (0.3–2.4)	.67
Any ischemic infarction after treatment	3.7 (1.4–10.7)	.007

Note:—IPH indicates intraparenchymal hemorrhage.