

On-line Table: Indications, aneurysm characteristics, and treatment techniques of 19 aneurysms treated in 18 patients with the Pipeline flow diverter

Patient ID	Indication				Location				Size				Technique			No. of PEDs							
	Inci-dental	Head-aches	Mass Effect	Recan	Cavernous	Para-ophthalmic	Sup Hypophyseal	PcomA	BA	Aneurysm (mm)	Neck (mm)	Volume (cm ³)	Fusiform	PED + Coils	PED Only	Packing Density	1	2	3	4	5		
1	1						1			18	7.4	1.25		1		13.9%	1					1	
1	1						1			14	7.6	0.83		1		12.4%	1					1	
2		1				1				17	6.3	1.07		1		7.1%	1					1	
3		1				1				12	4.5	0.38		1		23.3%	1					1	
4			1		1					28	13	0.84	1	1		5.7%						1	
5		1						1		10	5.1	0.38		1		26.8%	1					1	
6		1				1				9	4	0.13		1		14.3%	1					1	
7			1				1			16	6.6	1.63		1		7.2%	1					1	
8		1					1			12	4.8	0.34		1		21.8%	1					1	
9		1						1		20	9.3	2.43	1	1		1.4%	1					1	
10			1							18	9.8	4.26			1	0.0%						1	
11		1								10	7.5	0.62		1		0.0%	1					1	
12		1					1			7	5	0.22		1		0.0%	1					1	
13			1				1			30	8.7	3.60		1		0.0%	1					1	
14		1								5	4	0.05		1		0.0%	1					1	
15		1								8	5.3	0.26		1		0.0%	1					1	
16			1					1		35	9.1	3.97		1		0.0%	1					1	
17			1							6	5	0.04		1		0.0%	1					1	
18		1								26	18	1.64	1	1		0.0%	1					1	
Total	2	10	6	1	2	7	6	3	1	15.8	7.42	1.26	3	10	9	13.4%	7	7	3	1	1	1	
Mean																							