

On-line Table. Baseline clinical data, intervention, clinical, and angiographic outcomes

No.	Occlusion Site	Cause	Adjunctive Medication	Recapture, Period from Deployment to Attempt (min)	Baseline TIMI	Post-Tx TIMI	Baseline NIHSS	Discharge NIHSS
1	ICA terminus to A1 and M1, left	Cardioembolic	UK, 100,000 U; tirofiban, 0.5 mg	Failed (180)	0	3	22	0
2	ICA terminus to A1 and M1, left	Cardioembolic	UK, 100,000 U; tirofiban, 0.5 mg	Failed (120)	0	3	23	20
3	ICA terminus to M1, left	Traumatic dissection	UK, 100,000 U; tirofiban, 0.5 mg	No attempt	0	3	3→10	1
4	ICA terminus to A1 and M1, right	Cardioembolic	UK, 100,000 U; tirofiban, 1 mg	Failed (150)	0	3	18	3
5	ICA terminus to A1 and M1, right	Cardioembolic	UK, 200,000 U	Successful (20)	0	2	13	0
6	BA terminus	Cardioembolic	UK, 100,000 U; tirofiban 1 mg	Successful (30)	0	2	14	0
7	M1 to M2, right	Cardioembolic	UK, 100,000 U; tirofiban, 1 mg	Failed (120)	0	2	21	15
8 ^a	M1, right	Embolic, ICA to MCA	UK, 300,000 U	Successful (15)	0	2	15	0
9	M1 to M2, right	Cardioembolic	Tirofiban, 0.5 mg	No attempt	0	2	17	3

^aThis case was published as Suh et al.⁹