

On-line Table: Clinical and radiologic characteristics of the patient cohort

	All Patients (n = 55) (%)	Diffuse SAH (n = 33) (%)	Perimes SAH (n = 11) (%)	Periph SAH (n = 11) (%)
Sex				
Male	28 (51)	20 (61)	7 (64)	1 (9)
Female	27 (49)	13 (39)	4 (36)	10 (91)
Family history of aneurysms				
Yes	1 (2)	0	1 (9)	0
No	54 (98)	33 (100)	10 (91)	11 (100)
Smoking history				
Yes	21 (38)	12 (36)	5 (45)	4 (36)
No	34 (62)	21 (64)	6 (55)	7 (64)
Admission Hunt and Hess Scale				
1	17 (31)	6 (18)	8 (73)	3 (27)
2	19 (35)	12 (37)	3 (27)	4 (36)
3	14 (25)	10 (30)	0	4 (36)
4	5 (9)	5 (15)	0	0
Admission Fisher Scale				
1	0	0	0	0
2	10 (18)	0	1 (9)	9 (82)
3	30 (55)	18 (55)	10 (91)	2 (18)
4	15 (27)	15 (45)	0	0
Endovascular vasospasm treatment				
Yes	1 (2)	1 (3)	0	0
No	54 (98)	32 (97)	11 (100)	11 (100)
ICU length of stay, days				
0–1	7 (13)	1 (3)	2 (18)	4 (36)
2–7	23 (42)	11 (33)	7 (64)	5 (46)
≥8	25 (45)	21 (64)	2 (18)	2 (18)
Hospital length of stay, days				
2–7	20 (36)	6 (18)	6 (55)	8 (73)
8–14	22 (40)	14 (42)	5 (45)	3 (27)
≥15	13 (24)	13 (40)	0	0

Note:—Perimes indicates perimesencephalic; Periph, peripheral sulcal; ICU, intensive care unit.