

On-line Table: Most commonly selected responses for 14 scenarios and degree of agreement for the most commonly selected response^a

Scenario	Respondents Grouped by Years in Practice														
	All Respondents			Fellow			Attending <10 Years			Attending ≥10 Years			Respondents Grouped by Practice Type		
	Response	Agreement (%)		Response	Agreement (%)		Response	Agreement (%)		Response	Agreement (%)		Response	Agreement (%)	
8-mm 12F	D	43		D	41		D	47		D	40		D	43	
8-mm 30F	B	39		B	33		B	38		B	44		B	40	
8-mm 30M	B	38		B	41		D	36		B	42		B	39	
8-mm 60F	B	51		B	52		B	47		B	56		B	53	
10-mm 12F	D	56		D	56		D	57		D	56		D	57	
10-mm 30F	D	53		D	44		D	52		D	58		D	52	
10-mm 30M	D	53		D	44		D	52		D	58		D	51	
10-mm 60F	B	36		B	37		D	37		B	36		B	38	
15-mm 12F	D	67		D	73		D	68		D	63		D	67	
15-mm 30F	D	75		D	74		D	73		D	78		D	73	
15-mm 30M	D	75		D	74		D	72		D	80		D	73	
15-mm 60F	D	64		D	70		D	63		D	62		D	62	
8-mm lung cancer history	B	39		B	37		B	35		B	48		B	42	
10-mm multiple nodules	B	36		B	48		B	36		D	38		B	39	
Mean agreement		52			52			51			54			52	

Note:—B indicates report only in body; C, report in impression without recommendation; D, recommend ultrasound; M, male; F, female.

^a Differences in the most commonly selected response compared with radiologists in other groups.