

On-line Table 1: Clinical and angiographic data of 11 patients with bifurcation aneurysms treated with kissing-Y or crossing-Y stent-assisted coiling

Classification	Anterior Circulation (Nos. 1–6)						Posterior Circulation (Nos. 7–11)				
	1	2	3	4	5	6	7	8	9	10	11
Case and No.											
Age (yr)/sex	58/M	42/F	73/M	64/M	52/M	53/F	56/F	53/F	78/F	67/F	68/F
Presentation	SAH	Dizz	Dizz	SAH	Ha	SAH	Inc	Inc	SAH	Ha	Inc
HH scale	2	0	0	2	0	2	0	0	2	0	0
Bifurcation site	AcomA	AcomA	MCA-Bif	MCA-Bif	MCA-Bif	PcomA-fetal PCA	BA-Bif	BA-Bif	BA-Bif	BA-Bif	BA-Bif
Wide-neck	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Size	Small	Small	Small	Small	Small	Small	Small	Small	Small	Large	Small
Stent strategy	Planned	Non-P	Non-P	Non-P	Non-P	Non-P	Non-P	Planned	Planned	Planned	Planned
Y-Stent	Crossing	Crossing	Crossing	Crossing	Crossing	Crossing	Crossing	Crossing	Crossing	Kissing	Kissing
1st Stent	EP	EP	NF	NF	EP	EP	EP	EP	EP	NF	EP
	4.5/22	4.5/22	4.5/15	2.5/15	4.5/22	4.5/22	4.5/28	4.5/28	4.5/28	3.5/30	4.5/28
2nd Stent	EP	EP	NF	NF	EP	EP	EP	EP	EP	EP	EP
	4.5/14	4.5/22	4.5/15	2.5/15	4.5/22	4.5/22	4.5/28	4.5/22	4.5/22	4.5/28	4.5/28
Catheterization difficulty	No	No	Yes	Yes	No	No	No	No	No	Yes	Yes
Complications	No	Thrombus	No	No	No	No	No	No	No	No	No
Initial results	Class I	Class I	Class II	Class II	Class II	Class I	Class I	Class I	Class I	Class III	Class II
mRS/discharge	1	1	0	1	0	1	1	0	1	0	0
Angiographic FU interval (mo)	6	6	11	15	6	8	12	7	7	24	7
Results	Cured	Cured	Cured	Cured	Cured	Cured	Cured	Cured	Cured	Rec ^a	Stable
Retreatment	0	0	0	0	0	0	0	0	0	1	0
Clinical FU											
Interval (mo)	6	6	18	28	10	13	12	7	7	24	20
mRS/follow-up	1	1	0	1	0	1	1	0	1	0	0

Note:—Dizz indicates dizziness; Inc, incidental; Ha, headache; Non-P, non-planned (salvaging); EP, Enterprise; NF, Neuroform; Rec, recanalization; FU, follow-up; HH scale, Hunt and Hess scale.

^aRecanalization for coil compaction and retreatment once a complete occlusion was obtained.