

On-line Table 1: CTP parameters in patients who achieved LRC with RT compared with LRF^a

		Percentage Change from Pre-RT to			
LRC Status		Week 2, LRF (n = 2); LRC (n = 13)	Week 4, LRF (n = 2); LRC (n = 12)	Week 6, LRF (n = 1); LRC (n = 10)	6 Weeks Post-RT, LRF (n = 2); LRC (n = 10)
BF	LRF	-18.1 (15.7)	315.7 (470.8)	-69.0 (NA)	-46.2 (40.7)
	LRC	27.5 (47.7)	-5.3 (60.3)	31.8 (90.1)	-29.1 (49.1)
	P value	.046	.511	NA	.663
BV	LRF	-18.8 (10.9)	30.5 (81.3)	-61.5 (NA)	-38.8 (43.9)
	LRC	13.3 (42.7)	-18.4 (32.4)	5.5 (55.8)	-25.0 (38.7)
	P value	.053	.549	NA	.737
MTT	LRF	10.7 (9.4)	-37.6 (47.5)	41.7 (NA)	21.6 (3.4)
	LRC	-3.8 (32.7)	21.4 (106.0)	-1.5 (55.2)	18.4 (45.8)
	P value	.245	.281	NA	.834
CP	LRF	-21.3 (48.2)	360.6 (509.2)	-28.1 (NA)	53.5 (166.3)
	LRC	131.1 (289.3)	105.3 (288.4)	16.2 (165.7)	80.7 (283.5)
	P value	.106	.606	NA	.868

^aData are presented as mean ± SD. P values are calculated from an independent samples t test comparing mean percentage change in the LRC versus LRF groups.

On-line Table 2: BF (mL/100 g/min) in patients who achieved LRC with RT compared with LRF

	Patient No.	Baseline	Week 2	Week 4	Week 6	6 Weeks Post-RT
LRF	1	52.1	36.9	390.1	16.2	43.1
	2	54.7	50.9	45.3		13.7
LRC	1	95.2	81.3	34.8	167.2	103.0
	2	161.3	259.5	162.3	63	131.2
	3	48.3	87.5	84.4	50.8	22.5
	4	96.4	158.6	145.6	271.8	178.7
	5	76.6	113.0	94.8	58.7	24.4
	6	28.3	55.7	57.0	58.4	16.9
	7	119.6	115.8	54.3	92.8	99.0
	8	85.3	113.0	75.5	83.2	30.6
	9	174.6	181.3	31.5	23.9	21.1
	10	208.2	92.0	41.8	508.2	
	11	134	237.4	143.5		87.6
	12	252.6	230.4	174.1		
	13	53.7	40.6			