

**On-line Table: Baseline characteristics of the patients<sup>a</sup>**

Outcome	Entire Cohort (N = 202)	Studied Cohort (n = 196)	SPAN-100-Positive (n = 15)	SPAN-100-Negative (n = 181)
Median age (yr, range)	72 (25, 86)	72 (25, 86)	80 (76, 86)	70 (25, 84)
Age categories (n/N)				
<65 yr	27% (54/202)	27% (53/196)	0% (0/15)	29% (53/181)
65–79 yr	59% (119/202)	59% (115/196)	40% (6/15)	60% (109/181)
≥80 yr	14% (29/202)	14% (28/196)	60% (9/15)	10% (19/181)
Sex, female (n/N)	60% (122/202)	60% (117/196)	67% (10/15)	59% (107/181)
Median NIHSS score (range)	17 (8, 26)	17 (8, 26)	23 (18, 26)	16 (8, 26)
Median ASPECTS (range)	8 (3, 10)	8 (3, 10)	8 (5, 9)	8 (3–10)
ASPECTS categories (n/N)				
≤5	13% (25/199)	12% (24/193)	14% (2/14)	12% (22/179)
6–10	87% (174/199)	88% (169/193)	86% (12/14)	88% (157/179)
Hemispheric occlusion side (left) (n/N)	51% (103/202)	50% (98/196)	80% (12/15)	48% (86/181)
Prestroke mRS (n/N)				
0	62% (125/202)	64% (125/196)	67% (10/15)	64% (115/181)
1	15% (30/202)	15% (30/196)	20% (3/15)	15% (27/181)
2	11% (22/202)	11% (22/196)	7% (1/15)	12% (21/181)
3	0% (1/202)	1% (1/196)	0% (0/15)	1% (1/181)
Not determined	12% (24/202)	9% (18/196)	7% (1/15)	9% (17/181)
Median systolic blood pressure (mm Hg) (range)	146 (99, 220)	146 (99, 220)	154 (115, 208)	145 (99–220)
Median time from stroke onset to groin puncture (min) (range)	238 (72, 714)	239 (72, 714)	280 (140, 450)	235 (72–714)
Median time from stroke onset to TIC1 2b or 3 (or final DSA run) (min) (range)	282 (100, 800)	287 (100, 800)	322 (188, 483)	282 (100–800)
Stroke onset to successful revascularization categories (n/N)				
0–4.5 hr	46% (92/201)	45% (87/195)	20% (3/15)	47% (84/180)
4.5–8 hr	50% (101/201)	51% (100/195)	73% (11/15)	49% (89/180)
≥8 hr	4% (8/201)	4% (8/195)	7% (1/15)	4% (7/180)
IV-tPA administered (vs contraindicated or bridging) (n/N)	37% (75/202)	37% (73/196)	53% (8/15)	36% (65/181)
Medical history				
Diabetes mellitus	15% (30/202)	15% (30/196)	40% (6/15)	13% (24/181)
Myocardial disease	18% (36/202)	18% (35/196)	27% (4/15)	17% (31/181)
Systemic hypertension	63% (127/202)	64% (126/196)	80% (12/15)	63% (114/181)
Atrial fibrillation	38% (77/202)	39% (77/196)	60% (9/15)	38% (68/181)
Hyperlipidemia	39% (79/202)	39% (77/196)	40% (6/15)	39% (71/181)
Smoker (past or current)	19% (39/202)	19% (38/196)	7% (1/15)	20% (37/181)
Previous ischemic stroke	10% (21/202)	10% (19/196)	7% (1/15)	10% (18/181)
Previous hemorrhagic stroke	1% (3/202)	2% (3/196)	0% (0/15)	2% (3/181)
Occlusion location				
ICAT				
M1	18% (36/196)	18% (34/190)	20% (3/15)	18% (31/175)
M2	66% (130/196)	66% (126/190)	53% (8/15)	67% (118/175)
M3	14% (27/196)	14% (27/190)	27% (4/15)	13% (23/175)
M3	2% (3/196)	2% (3/190)	0% (0/15)	2% (3/175)

**Note:**—ICAT indicates intracranial carotid T-type occlusion; n/N, group number/entire cohort number.

<sup>a</sup> Six patients had missing NIHSS scores at baseline and the most proximal occlusion location.