

Supplemental Online Content

Zatoński WA, Zatoński M, Janik-Koncewicz K, Wojtyła A. Alcohol-related deaths in Poland during a period of weakening alcohol control measures. *JAMA*. Published March 16, 2021. doi:10.1001/jama.2020.25879

eTable. List of Causes of Death Wholly Attributable to Alcohol Consumption Included in the Analysis

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable. List of Causes of Death Wholly Attributable to Alcohol Consumption Included in the Analysis

Cause of death	<i>ICD-10</i> code
Mental and behavioural disorders due to use of alcohol	F.10.x
Alcohol intoxication	F10.0
Harmful use of alcohol	F10.1
Alcohol dependence	F10.2
Alcohol withdrawal state	F10.3
Alcohol withdrawal state with delirium	F10.4
Alcohol-induced psychotic disorder	F10.5
Alcohol-induced amnestic disorder	F10.6
Alcohol-induced residual and late onset psychotic disorder	F10.7
Poisoning due to alcohol	
Accidental poisoning by and exposure to alcohol	X45
Intentional self-poisoning by and exposure to alcohol	X65
Poisoning by and exposure to alcohol, undetermined intent	Y15
Liver disorders due to alcohol	K70.x
Alcoholic fatty liver	K70.0
Alcoholic hepatitis	K70.1
Alcoholic fibrosis and sclerosis of liver	K70.2
Alcoholic cirrhosis of liver	K70.3
Alcoholic hepatic failure	K70.4
Alcoholic liver disease, unspecified	K70.9
Gastrointestinal disorders due to alcohol	
Alcoholic gastritis	K29.2
Alcohol-induced acute pancreatitis	K85.2
Alcohol-induced chronic pancreatitis	K86.0
Other disorders due to alcohol	
Alcohol-induced pseudo-Cushing's syndrome	E24.4
Degeneration of nervous system due to alcohol	G31.2
Alcoholic polyneuropathy	G62.1
Alcoholic myopathy	G72.1
Alcoholic cardiomyopathy	I42.6

Sources:

1. Rehm J, Baliunas D, Borges GLG, et al. The relation between different dimensions of alcohol consumption and burden of disease – an overview. *Addiction* 2010; 105(5): 817-843.
2. Roberts E, Morse R, Epstein S, et al. The prevalence of wholly attributable alcohol conditions in the United Kingdom hospital system: a systematic review, meta-analysis and meta-regression. *Addiction* 2019; 114(10): 1726-1737.

Notes:

1. Rehm et al. included in their list six additional health conditions wholly attributable to alcohol: foetal alcohol syndrome(Q86.0), foetus and newborn affected by maternal use of alcohol (P04.3), maternal care for (suspected) damage to foetus from alcohol (O.35.4), excess blood alcohol (R78.0), toxic effect of alcohol (T51), evidence of alcohol involvement determined by blood alcohol level (Y90). In our analysis these six conditions were excluded.
2. Data on alcohol-induced pseudo-Cushing's syndrome (E24.4) for both men and women, and on alcoholic myopathy for women were not available in Polish mortality database.