

Supplementary Table SIV Interaction of total T with diagnostic group with respect to NEFA parameters

Dependent variables	Independent variables	β coefficient \pm SE	P-value	R ²
NEFA ₀ (μ mol/l)	Diagnostic group \times total T	-0.002 ± 0.001	0.073	13.9
	Total T	0.002 ± 0.001	0.15	
	Diagnostic group	-0.038 ± 0.055	0.486	
g _s (mmol/l)	Diagnostic group \times total T	0.002 ± 0.002	0.135	4.7
	Total T	-0.002 ± 0.002	0.157	
	Diagnostic group	0.134 ± 0.065	0.0449	
log K _C (%/min)	Diagnostic group \times total T	-0.003 ± 0.002	0.148	4.9
	Total T	0.003 ± 0.002	0.126	
	Diagnostic group	-0.207 ± 0.080	0.0127	
Latency (τ) (min)	Diagnostic group \times total T	-0.002 ± 0.002	0.359	4.4
	Total T	0.002 ± 0.002	0.275	
	Diagnostic group	-0.084 ± 0.071	0.2408	
log Φ (mmol/l)	Diagnostic group \times total T	-0.007 ± 0.003	0.026	10.3
	Total T	0.007 ± 0.003	0.036	
	Diagnostic group	-0.287 ± 0.137	0.0422	
S _{FFA} (μ mol/l/min)	Diagnostic group \times total T	0.001 ± 0.002	0.965	2.5
	Total T	-0.001 ± 0.002	0.827	
	Diagnostic group	0.114 ± 0.071	0.1137	
K _{FFA} (%/min)	Diagnostic group \times total T	-0.001 ± 0.002	0.761	5.7
	Total T	0.001 ± 0.002	0.57	
	Diagnostic group	-0.027 ± 0.065	0.6739	